

THE SCIENCE OF SMILES

Neuroscience-Driven Physiotherapy Strategies for Bell's Palsy

DR. DARSHPREET KAUR (PT)

DR. (PROF.) GUNJAN KUMAR



BlueRoseONE^{.com}
S t o r i e s M a t t e r

New Delhi • London

BLUEROSE PUBLISHERS
India | U.K.

Copyright © Dr. Darshpreet Kaur, Dr. (Prof.) Gunjan Kumar 2026

All rights reserved by author. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the author. Although every precaution has been taken to verify the accuracy of the information contained herein, the publisher assumes no responsibility for any errors or omissions. No liability is assumed for damages that may result from the use of information contained within.

BlueRose Publishers takes no responsibility for any damages, losses, or liabilities that may arise from the use or misuse of the information, products, or services provided in this publication.



For permissions requests or inquiries regarding this publication,
please contact:

BLUEROSE PUBLISHERS
www.BlueRoseONE.com
info@bluerosepublishers.com
+91 8882 898 898
+4407342408967

ISBN: 978-93-7542-000-2

Cover Design: Deepak Katheria
Typesetting: Manisha Rawat

First Edition: May 2026

Dedication

Dedicated to all patients who trusted us for treatment of Bell's palsy especially during testing times of COVID; and to clinicians & physiotherapists committed to restoring function, confidence, and dignity through evidence-based care.

Acknowledgements

The authors gratefully acknowledge the multidisciplinary teams, patients, and institutions whose collaboration, trust, and commitment contributed to the clinical data and rehabilitation protocols presented in this book. Their collective engagement enabled the translation of clinical practice into structured, evidence-informed approaches.

The authors extend particular appreciation to the team at **Bihar Neuro Diagnostic Centre** and **Neuro Vihar, Rukanpura, Patna**, whose sustained professionalism, discipline, and dedication were fundamental to the successful execution of this work. The consistent efforts of this team created a robust clinical environment that supported both patient care and academic rigor.

Special recognition is accorded to the neurotechnicians—**Mr. Deepak, Mr. Aditya, Mr. Suraj, Ms. Manisha Mohanty, Mr. Luvkush, Mr. Babin, Mr. Rabiul, and Mr. Aman**—for their technical expertise, precision, and steadfast contribution to diagnostic and monitoring processes that informed clinical decision-making.

The authors also acknowledge, with sincere appreciation, the physiotherapy team for their clinical excellence, collaborative spirit, and commitment to rehabilitation outcomes: **Dr. Ravi (PT), Dr. Umakant (PT), Dr. Tarique Aziz (PT), Dr. Sakshi (PT), Dr. Ayushi (PT), Dr. Om Prakash (PT), Dr. Nitish (PT), Dr. Bibhuti Jha (PT), Dr. Kriti (PT), Dr. Rupa Shandilya (PT), and Dr. Prince Yadav (PT)**. Their contributions exemplify high standards of professional practice and patient-centred rehabilitation.

Contributors

Dr.(Prof) Ajay Kumar Singh

Director Cum Founder Bihar Neurodiagnostic Centre

Ex. Director, Dean, Vice-chancellor, HOD of Neurology, IGIMS,
Patna

Ex. Associate Professor, NIMS, Hyderabad

Dr. (Prof) Gunjan Kumar

Professor, Dept. of Neurology, PMCH, Patna

Founder & Head, Dept. of Neurology, Neuro Vihar

Founder HOD Neurology AIIMS-Patna

Dr. Sarsij Sharma

Additional Professor in Plastic Surgery, AIIMS-Patna

Dr. Amrita Kumari

Associate Professor, Dept. of Anatomy, PMCH, Patna

Dr. Vihag Raman

Director & Sr. Consultant Radiologist, Raman Imaging, Patna

Ex. Registrar, Manipal Hospital, Bangalore

Dr. Nidhi Billore (PT)

Ex. Assistant Professor, RV College of Physiotherapy, Bangalore

Certified Yoga Instructor, South Korea

Dr. Vidhi Singh (PT)

Research Scholar, Jayoti Vidhyapeeth Women's University, Jaipur

Dr. Darshpreet Kaur(PT)

Director- Physiotherapy, Institute of Neurosciences, Jeevandan Health,
Danapur.

Research Scholar, Jayoti Vidhyapeeth Women's University, Jaipur

Director- Dept. of Physiotherapy, Neuro Vihar Group of Centres

Dr. (Prof) Sneha Vishwanil Katke (PhD, MPT)

Principal cum Professor Bharati Vidyapeeth School of Physiotherapy,
Sangli, Maharashtra.

Foreword

Facial palsy presents not only as a neurological deficit but as a profound functional and psychosocial challenge. This book aims to bridge anatomy, diagnostics, and rehabilitation into a cohesive clinical framework, supported by real-world data and structured protocols.

Preface

Facial palsy represents a complex neurological condition that extends far beyond a visible motor deficit. Effective management requires a nuanced understanding of facial neuroanatomy, underlying pathology, diagnostic interpretation, and the principles of neurorehabilitation. While significant advances have been made in imaging modalities, pharmacological interventions, and surgical techniques, **structured, protocol-driven rehabilitation continues to remain underrepresented in mainstream clinical literature.**

This book has been developed to address this critical gap. It integrates **electrodiagnostic evaluation, physiotherapy-driven rehabilitation strategies, and outcome-based proprietary protocols** derived from extensive clinical experience in managing patients with facial nerve disorders. Emphasis is placed on translating diagnostic findings into targeted rehabilitation interventions, thereby bridging the divide between investigation and functional recovery.

The content is grounded in contemporary neuroscience and rehabilitation science, with practical insights drawn from real-world clinical practice. Special attention is given to objective assessment tools, staged rehabilitation planning, neuromuscular re-education, and functional outcome monitoring—areas that are often fragmented or inadequately addressed in conventional texts.

This book is intended for **neurologists, physiotherapists and other rehabilitation professionals** who seek a comprehensive, evidence-informed, and clinically applicable guide to facial palsy management. By presenting an integrated, multidisciplinary approach, it aims to support improved clinical decision-making and optimize patient-centred outcomes.

Dr. (Prof) Gunjan Kumar

Dr. Darshpreet Kaur (PT)

Foreword by Dr. Mansukhani



Bombay Hospital & Medical Research Centre
DEPARTMENT OF CLINICAL NEUROPHYSIOLOGY
EMG & EVOKED POTENTIALS

12, New Marine Lines, Mumbai 400 020. • Tel: 2206 76 76 Ext: 374 • Fax: 2208 08 71



Bell's palsy remains one of the most common causes of acute facial paralysis, presenting not only with functional impairment but also with profound psychological and social consequences for those affected. The sudden onset of facial weakness challenges patients and clinicians alike, demanding timely, evidence-based, and compassionate care. Among the various management approaches, physiotherapy plays a pivotal role in promoting recovery, preventing secondary complications, and restoring confidence and quality of life.

This book offers a comprehensive and clinically relevant exploration of the physiotherapy management of Bell's palsy. It bridges the gap between foundational neurophysiological principles and practical therapeutic application, providing readers with a clear understanding of facial nerve anatomy, pathophysiology, assessment strategies, and stage-wise rehabilitation protocols. Emphasis is placed on individualized care, recognizing that recovery from Bell's palsy varies widely and requires thoughtful clinical reasoning rather than a one-size-fits-all approach.

What distinguishes this work is its focus on functional recovery through safe and progressive physiotherapy interventions. The text highlights the importance of early intervention, patient education, facial muscle re-education, neuromuscular control, and the prevention of synkinesis and long-term dysfunction. By integrating current evidence with clinical experience, the book equips physiotherapists, students, and rehabilitation professionals with practical tools to optimize outcomes.

Equally important is the attention given to the emotional and psychosocial impact of facial paralysis. Physiotherapists are often closely involved in long-term patient care, placing them in a unique position to support, motivate, and guide individuals through what can be a distressing recovery journey. This book reinforces the role of the therapist not only as a clinician, but also as an educator and advocate for patient-centered care.

I commend DR. DARSHPREET KAUR (PT) for addressing a topic of significant clinical importance with clarity and depth and provide a **proprietary and outcome-based** book. This book will serve as a valuable resource for undergraduate and postgraduate students, practicing physiotherapists, and other healthcare professionals involved in the multidisciplinary management of Bell's palsy. It is my hope that this work will contribute meaningfully to improved rehabilitation practices and better patient outcomes.

Dr. K.A.MANSUKHANI
DEPARTMENT HEAD

Foreword by Dr.Pravin



I'm delighted to introduce and write the foreword for "The Science of Smiles: Modern Physiotherapy Approaches in Bell's Palsy". This book is a must-have for clinicians and students seeking to enhance their understanding of facial nerve disorders and evidence-based rehabilitation strategies. The authors have expertly distilled complex concepts into practical, clinically relevant insights, making it an invaluable resource for anyone working with patients with Bell's Palsy.

As a teacher, I'm always on the lookout for resources that can help my students and colleagues stay ahead in the game, and this book delivers. Bell's Palsy is a condition that can be tricky to tackle, but Darsh Gunjan's work is a game-changer. Over the past five years, they've poured their heart and soul into developing a proprietary treatment protocol that's yielded impressive results with over 200 patients. And now, they're sharing their expertise with the world.

What sets this book apart is its comprehensive approach, combining facial nerve anatomy, electrodiagnostic evaluation, and rehabilitation protocols in one place. Whether you're a student, clinician, or seasoned pro, you'll appreciate the clarity and practicality of the content.

I'm excited to see the impact this book will have on the field of physiotherapy, and I'm confident it will become a go-to resource for anyone working with Bell's Palsy patients. Kudos to Dr. Darshpreet (PT) and Dr. Gunjan (DM-Neurology) for this outstanding contribution.

A handwritten signature in black ink, appearing to read 'Pravin'.

Pravin Kumar
Head,

Department of Physiotherapy,
National Institute For Locomotor Disabilities (Divyangjan),
Bon Hooghly, BT Road,
Kolkata, West Bengal, India.

Foreword By Dr. Jitendra

डॉ. जितेंद्र शर्मा
DR. JITENDRA SHARMA
निदेशक / Director
टेलीफोन / Telephone : 011-23232403
ईमेल / E-mail : diriph@nic.in



पंडित दीनदयाल उपाध्याय राष्ट्रीय शारीरिक दिव्यांगजन संस्थान
दिव्यांगजन सशक्तिकरण विभाग
सामाजिक न्याय और अधिकारिता मंत्रालय, भारत सरकार

Pt. Deendayal Upadhyaya National Institute
for Persons with Physical Disabilities (Divyangjan)
Department of Empowerment of Persons with Disabilities (Divyangjan)
Ministry of Social Justice & Empowerment, Government of India

File No. 01-01/2024 - Director Secretariat / 12 / 2026

05th February '26



When I first encountered this manuscript, the first thought that came to my mind was that this book speaks with authority yet never condescends, that delves deep yet remains accessible, that acknowledges complexity while offering clarity.


Bell's Palsy, though common enough that most neurologists see cases regularly, remains a condition shrouded in uncertainty for those who experience it. The sudden paralysis of one side of the face—often discovered upon waking or within hours—carries with it a cascade of questions, fears, and practical challenges that extend far beyond the purely medical. How long will this last? Will my face return to normal? What can I do to help my recovery? These questions deserve answers that are both scientifically sound and genuinely useful.

This is precisely what the authors have achieved. The collaboration between a neurologist and a physiotherapist has produced something greater than the sum of its parts. Where the neurologist brings deep understanding of the underlying pathophysiology; Diagnostic nuances, and evidence-based treatment protocols, the physiotherapist contributes practical wisdom about rehabilitation, facial exercises, and the daily realities of recovery. Together, they have woven these perspectives into a seamless whole.

What impressed me most was their commitment to clarity without oversimplification. They explain the anatomy of the facial nerve in terms anyone can grasp, yet they don't shy away from discussing the subtleties of differential diagnosis or the limitations of current treatment options. They present rehabilitation exercises with step-by-step precision, while also acknowledging that recovery timelines vary and that patience must accompany effort.

The result is a book that will serve multiple audiences well: patients seeking to understand and actively participate in their recovery, family members wanting to provide informed support, and healthcare professionals looking for a comprehensive yet approachable resource to share with those they treat.

In an age when medical information is abundant but wisdom is scarce, this book offers both. I commend it to anyone touched by Bell's Palsy, confident that they will find within these pages not just information, but genuine guidance for the journey ahead



Dr. Jitendra Sharma (PT)

Contents


Chapter 1: Overview Of Facial Palsy	1
Chapter 2: Anatomy Of The Facial Nerve.....	6
Chapter 3: Neurological Aspects Of Bell's Palsy.....	16
Chapter 4: Clinical Evaluation And Grading Systems.....	22
Chapter 5: Radiological Investigations In Facial Palsy	29
Chapter 6: Electrophysiological Tests Used In Facial Palsy	35
Chapter 7: Pharmacological And Acute Phase Management Of Facial Palsy.....	43
Chapter 8: Plastic And Reconstructive Surgical Approaches In Facial Palsy.....	49
Chapter 9: Physiotherapy And Rehabilitation Techniques In Facial Palsy.....	56
Chapter 10: Proprietary Rehabilitation Protocol And Clinical Data	66
Chapter 11: Multidisciplinary Approach And Future Directions In Facial Palsy Management.....	73
Chapter 12: Evidence Based Practices.....	78
About The Authors	91

CHAPTER 1

Overview of Facial Palsy

Dr. Darshpreet Kaur (PT)

Introduction



Facial palsy represents a spectrum of conditions characterized by weakness or paralysis of the muscles of facial expression due to dysfunction of the facial nerve (cranial nerve VII). Among these, Bell's palsy—also known as idiopathic facial palsy—is the most common cause of acute peripheral facial paralysis encountered in clinical practice. Understanding its presentation, pathophysiology, evaluation, and early management is fundamental for optimizing functional and cosmetic outcomes.

Bell's Palsy: Definition and Clinical Presentation

Bell's palsy is defined as an acute, idiopathic, lower motor neuron paralysis of the facial nerve, typically affecting one side of the face. The hallmark feature is a **sudden onset of unilateral facial weakness**, which usually progresses rapidly and reaches its maximum severity within 24 to 48 hours.

Clinically, patients present with facial asymmetry due to weakness of the muscles of facial expression on the affected side. Common features include inability to close the eye, flattening of the nasolabial fold, drooping of the angle of the mouth, difficulty in smiling, and impaired facial expressions. Because the facial nerve also carries parasympathetic and sensory fibers, patients may experience altered lacrimation, hyperacusis, and disturbances in taste sensation over the anterior two-thirds of the tongue.

Etiology and Pathophysiology

Bell's palsy is caused by **inflammation and edema of the facial nerve**, particularly within the narrow confines of the facial canal. This swelling leads to compression, ischemia, and conduction block of the nerve fibers.

Although the exact etiology remains unclear, viral reactivation—most notably **herpes simplex virus type 1 (HSV-1)**—is widely considered a major contributing factor. Other conditions may also affect the facial nerve; however, these are typically associated with additional neurological or systemic signs, which are absent in classical Bell's palsy. The unilateral nature of the condition reflects the involvement of a single facial nerve, as each side of the face is innervated independently.

Assessment of Facial Paralysis Severity

Accurate assessment of facial nerve dysfunction is essential for diagnosis, prognostication, and monitoring recovery. Two grading systems are most commonly used in clinical practice:

- **House–Brackmann (H–B) Scale**, which provides a global grading of facial function from normal to total paralysis
- **Sunnybrook Facial Grading System**, which offers a more detailed regional and functional assessment, including resting symmetry, voluntary movement, and synkinesis

These scales help clinicians objectively document the extent of paralysis and track changes over time.

Historical Perspective: Sir Charles Bell



Source: https://en.wikipedia.org/wiki/Charles_Bell

Sir Charles Bell, a Scottish anatomist and surgeon, made seminal contributions to the understanding of the facial nerve. In 1829, he described the anatomy and functional role of the facial nerve in facial expression. His work laid the foundation for distinguishing facial nerve palsy from other neurological conditions, and the disorder later came to be known as **Bell's palsy** in recognition of his contributions.

Vulnerability of Facial Muscles

Facial muscles are uniquely vulnerable in facial nerve palsy due to their anatomical and functional characteristics. They are located in the **subcutaneous plane**, with minimal protective cushioning compared to deeper skeletal muscles. Additionally, facial muscles depend entirely on the integrity of a single motor nerve—the facial nerve—for voluntary movement.

When neural input is disrupted, these muscles rapidly lose tone and coordination, predisposing them to **contractures, synkinesis, and long-term disfigurement** if appropriate rehabilitation is delayed.


Facial Contractures and Their Prevention

Facial contractures refer to permanent shortening or tightening of facial muscles resulting from prolonged disuse, abnormal reinnervation, or

improper healing. These changes can lead to facial stiffness, asymmetry, and restricted movement.

Early initiation of physiotherapy, including facial muscle re-education and appropriate neuromuscular stimulation, plays a critical role in preventing contractures and preserving facial symmetry.

Bell's Phenomenon



Bell's phenomenon is a protective ocular reflex characterized by **upward and outward rotation of the eyeball** when the patient attempts to close the eyelids. Its presence is clinically significant, as it indicates weakness of the orbicularis oculi muscle and incomplete eye closure.

Recognition of Bell's phenomenon alerts clinicians to the need for meticulous eye care, including lubrication and protective measures, to prevent corneal injury.

Role of Electroneuronography (ENoG)

Electroneuronography (ENoG) is a valuable electrodiagnostic tool in the assessment of Bell's palsy. It measures facial nerve conduction and quantifies the degree of neural degeneration by comparing the affected side with the unaffected side.

Findings typically include reduced amplitude and prolonged latency on the affected side. ENoG assists in grading severity, estimating prognosis, and guiding clinical decision-making, particularly in moderate to severe cases.

Initial Management of Bell's Palsy

The cornerstone of Bell's palsy management is **early intervention**. Prompt initiation of pharmacological therapy—primarily corticosteroids—combined with supportive care significantly improves

recovery outcomes. Eye protection, patient education, and timely referral for rehabilitation are integral components of early management.

Chapter Summary

Bell's palsy is a common yet potentially disabling condition requiring early recognition and structured management. A thorough understanding of its clinical features, assessment tools, and early therapeutic strategies forms the foundation for effective rehabilitation and optimal functional recovery.

Further reading

Hohman MH, Hadlock TA. Etiology, diagnosis, and management of facial palsy: 2000 patients at a facial nerve center. Laryngoscope. 2014 Jul;124(7):E283-93.

Peitersen E. Bell's palsy: the spontaneous course of 2,500 peripheral facial nerve palsies of different etiologies. Acta Otolaryngol Suppl. 2002;(549):4-30.

Adour KK, Byl FM, Hilsinger RL, Kahn ZM, Sheldon MI. The true nature of Bell's palsy: analysis of 1,000 consecutive patients. Laryngoscope. 1978 May;88(5):787-801

CHAPTER 2


Anatomy of the Facial Nerve

Dr. Amrita Kumari

Introduction

The facial nerve (cranial nerve VII) is a complex mixed nerve with motor, sensory, and parasympathetic components. Its intricate anatomical course, long intracranial and intratemporal pathway, and passage through narrow bony canals make it particularly vulnerable to pathological processes. A thorough understanding of its anatomy is essential for accurate diagnosis, imaging interpretation, surgical intervention, and rehabilitation in facial nerve disorders, especially Bell's palsy.

Functional Components of the Facial Nerve



The facial nerve contains four major functional components:

1. **Special Visceral Efferent (SVE) Fibers**

These motor fibers arise from the facial motor nucleus in the pons and supply the muscles derived from the second pharyngeal (branchial) arch. This includes the muscles of facial expression, stapedius, posterior belly of digastric, and stylohyoid muscles.

2. **General Visceral Efferent (GVE) Fibers**

These are preganglionic parasympathetic fibers that supply the pterygopalatine and submandibular ganglia. Through these ganglia, they regulate lacrimation and salivation, particularly of the lacrimal, submandibular, and sublingual glands.


3. **Special Visceral Afferent (SVA) Fibers**

These fibers are responsible for taste sensation from the anterior two-thirds of the tongue (excluding vallate papillae). They project centrally to the upper part of the nucleus tractus solitarius.

4. **General Somatic Afferent (GSA) Fibers**

The cell bodies of these fibers are located in the geniculate ganglion. Their central processes terminate in the spinal nucleus of the trigeminal nerve, while peripheral processes supply sensation to parts of the external ear.

Intracranial Course of the Facial Nerve




The facial nerve emerges from the **anterolateral aspect of the brainstem at the pontomedullary junction**. It consists of two roots:

- A larger **motor root**
- A smaller **sensory root**, known as the *nervus intermedius*

These roots pass through the cerebellopontine angle and enter the **internal acoustic meatus**. Within the meatus, the motor root occupies the superior position, the sensory root lies centrally, and the vestibulocochlear nerve lies inferiorly. At the fundus of the internal acoustic meatus, the two roots unite to form a common trunk that enters the facial canal.

Facial Canal (Fallopian Canal) please remove the background of this picture



Source: https://en.wikipedia.org/wiki/Facial_canal


The facial canal is a bony passage within the petrous part of the temporal bone, extending from the internal acoustic meatus to the stylomastoid foramen. It is approximately 3 cm long, making it the **longest bony canal traversed by any nerve in the human body**.

Contents of the Facial Canal

- Facial nerve
- Greater petrosal nerve (parasympathetic fibers)
- Nerve to stapedius (motor supply to stapedius muscle)
- Chorda tympani (taste and parasympathetic fibers)
- Stylomastoid artery (vascular supply)

Clinical Significance

Knowledge of the facial canal anatomy is critical during middle ear and temporal bone surgeries, including cochlear implantation. Injury to the nerve within this canal can result in facial paralysis and long-term disability.



Source [Neuroanatomy, Geniculate Ganglion](#) Copyright © 2025, StatPearls Publishing LLC.

Intratemporal Course and Geniculate Ganglion

Within the facial canal, the nerve first runs laterally above the vestibule of the inner ear to reach the medial wall of the middle ear. It then makes a sharp posterior bend above the oval window known as the **external genu**, where the **geniculate ganglion** is located.

From this point, the nerve runs vertically downward along the posterior wall of the middle ear, behind the pyramid, and exits the skull through the **stylomastoid foramen**.

The geniculate ganglion serves as a sensory and parasympathetic relay station and is particularly vulnerable due to its location within a narrow bony segment, making it a common site of pathology in facial nerve disorders.

Extracranial Course and Parotid Plexus


After emerging from the stylomastoid foramen, the facial nerve gives off the **posterior auricular nerve**, supplying the occipital belly of occipitofrontalis and auricular muscles. It then winds around the styloid process and enters the parotid gland by piercing its posteromedial surface.

Within the parotid gland, the nerve divides into:

- **Temporofacial division**
- **Cervicofacial division**

These divisions form the **parotid plexus**, from which five terminal branches arise.

Terminal Branches of the Facial Nerve



The five terminal branches are:

1. **Temporal Branch**— supplies the frontal belly of occipitofrontalis, orbicularis oculi, and corrugator supercilii
2. **Zygomatic Branches (upper and lower)** – supply muscles around the eye, nose, and upper lip
3. **Buccal Branch** – supplies buccinator and muscles of the upper lip
4. **Marginal Mandibular Branch** – supplies muscles of the lower lip and chin, including risorius
5. **Cervical Branch** – supplies the platysma muscle

Role of Chorda Tympani

The chorda tympani is a mixed nerve branch of the facial nerve. It carries:

- **Taste sensation** from the anterior two-thirds of the tongue
- **Preganglionic parasympathetic fibers** to the submandibular ganglion

Postganglionic fibers from this ganglion supply the submandibular, sublingual, and anterior lingual glands, contributing to salivation.

Facial Nerve and Autonomic Functions

Lacrimation is mediated by parasympathetic fibers originating in the facial nerve and passing through the greater petrosal nerve to the pterygopalatine ganglion. Salivation is controlled through the chorda tympani and submandibular ganglion. Damage to these fibers can result in dry eye, reduced salivation, and taste disturbances.

Anatomical Basis of Bell's Palsy

The **labyrinthine segment** of the facial nerve is the most commonly affected segment in Bell's palsy. It is the narrowest portion of the facial canal and is highly susceptible to compression. Inflammation and edema within this confined space lead to ischemia, impaired conduction, and subsequent facial paralysis.

The narrow anatomy of the facial canal, particularly near the geniculate ganglion, plays a critical role in the pathophysiology of Bell's palsy.

Stylomastoid Foramen as a Clinical Landmark

The stylomastoid foramen marks the point where the facial nerve exits the skull and enters the parotid gland. Due to its narrow passage and anatomical relationships, this region is vulnerable to traumatic and surgical injury, making it an important landmark in facial nerve palsy.

Unilateral Nature of Bell's Palsy

Each facial nerve supplies the muscles of facial expression on the **ipsilateral side**. Since Bell's palsy involves pathology of a single facial nerve, the resulting paralysis is characteristically unilateral.

Supranuclear versus Infranuclear Lesions

- **Supranuclear (Upper Motor Neuron) Lesions**
Occur above the facial nucleus in the pons. They typically cause contralateral weakness of the lower two-thirds of the face while sparing the forehead due to bilateral cortical innervation.
- **Infranuclear (Lower Motor Neuron) Lesions**
Occur at or below the facial nucleus and result in complete ipsilateral facial paralysis, including the forehead. These lesions may also involve impaired lacrimation, hyperacusis, and taste disturbances.

Imaging Landmarks for Facial Nerve Evaluation

On MRI and CT imaging, the facial nerve can be traced using the following landmarks:

- Stylomastoid foramen
- Tympanomastoid suture
- Tragal cartilage
- Posterior belly of digastric muscle
- Peripheral facial nerve branches

These landmarks assist in localizing lesions and planning surgical interventions.

Anatomical Relations and Differential Diagnosis

Cranial nerves closely related to the facial nerve include:

- **Trigeminal nerve (CN V)**
- **Vestibulocochlear nerve (CN VIII)**
- **Nervus intermedius**
- **Glossopharyngeal nerve (CN IX)**

Pathologies at the cerebellopontine angle or internal acoustic meatus may simultaneously affect these nerves, leading to overlapping clinical features.

Surgical Decompression in Severe Bell's Palsy

In cases requiring surgical decompression, the facial nerve within the **fallopian canal**, particularly the **labyrinthine segment and region proximal to the geniculate ganglion**, must be carefully evaluated. These segments are the most common sites of compression and inflammation.

Chapter Summary

The facial nerve's complex anatomy and prolonged bony course explain both its diverse functions and its vulnerability to injury. A sound anatomical understanding is indispensable for clinicians managing facial nerve palsy, guiding diagnosis, imaging, surgical planning, and rehabilitation.

Further Reading:

Proctor B. The anatomy of the facial nerve. Otolaryngol Clin North Am. 1991 Jun;24(3):479-504. PMID: 1762772.

Guerrier Y. Le nerf facial. Quelques points d'anatomie topographique [The facial nerve. Points of topographic anatomy]. Ann Otolaryngol Chir Cervicofac. 1975 May-Jun;92(4-5):167-71. French. PMID: 1082736.

Donaldson JA, Anson BJ. Surgical anatomy of the facial nerve. Otolaryngol Clin North Am. 1974 Jun;7(2):289-308. PMID: 4599258.

Seneviratne SO, Patel BC. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): May 23, 2023. Facial Nerve Anatomy and Clinical Applications.

CHAPTER 3

Neurological Aspects of Bell's Palsy

Dr. (Prof) Ajay Kumar Singh
Dr. (Prof) Gunjan Kumar
Dr. (Prof) Sneha V. Ketke

Introduction

Facial palsy represents a neurological disorder resulting from dysfunction of the facial nerve or its central connections. From a neurological perspective, facial palsy is broadly classified based on the level of lesion into central (supranuclear) and peripheral (infranuclear) types. Accurate localization of the lesion is critical, as it determines etiology, prognosis, and management strategy. This chapter focuses on the neurological mechanisms, clinical differentiation, and common syndromes associated with facial palsy.

Neuroanatomical Basis of Facial Nerve Function

The facial nerve nucleus is located in the lower pons and receives bilateral cortical innervation for the upper facial muscles, while the lower facial muscles receive predominantly contralateral cortical input. This anatomical arrangement explains the distinct clinical patterns observed in central versus peripheral facial palsy.

The facial nerve also carries parasympathetic fibers responsible for lacrimation and salivation, as well as sensory fibers for taste. Therefore, neurological involvement of the facial nerve often presents with associated autonomic and sensory disturbances.

Central (Supranuclear) Facial Palsy

Central facial palsy results from lesions affecting the corticobulbar pathways above the facial nerve nucleus. Common causes include ischemic or hemorrhagic stroke, brain tumors, demyelinating disorders, and traumatic brain injury.

Clinically, central facial palsy is characterized by:

- Weakness of the **contralateral lower half of the face**
- **Preservation of forehead movements** due to bilateral cortical innervation
- Absence of taste disturbance, hyperacusis, or lacrimation abnormalities

Patients may present with accompanying neurological deficits such as hemiparesis, aphasia, or sensory loss, reflecting broader cerebral involvement.

Peripheral (Infranuclear) Facial Palsy

Peripheral facial palsy occurs due to damage to the facial nerve nucleus, its intrapontine fibers, or the nerve along its extracranial course. Bell's palsy is the most common example.

Clinical features include:

- Complete **ipsilateral facial paralysis**, including forehead and eyelid
- Loss of facial expression
- Possible impairment of lacrimation, salivation, and taste
- Hyperacusis due to stapedius muscle involvement

The extent of associated symptoms depends on the anatomical level of the lesion.

Bell's Palsy: Neurological Considerations

Bell's palsy is an acute idiopathic lower motor neuron facial palsy. Neurologically, it is believed to result from inflammatory edema of the facial nerve, most commonly affecting the labyrinthine segment within the facial canal.

Key neurological features include:

- Sudden onset facial weakness progressing over 24–48 hours
- Absence of central neurological signs
- Variable involvement of autonomic and sensory fibers

Early neurological assessment is essential to exclude secondary causes such as stroke, tumors, or infections.

Ramsay Hunt Syndrome

Ramsay Hunt syndrome results from reactivation of varicella-zoster virus in the geniculate ganglion. It represents a more severe form of peripheral facial palsy.

Neurological features include:

- Facial paralysis
- Severe otalgia
- Vesicular eruptions in the external auditory canal or pinna
- Hearing loss and vertigo due to vestibulocochlear nerve involvement

Prognosis is generally poorer than Bell's palsy, emphasizing the importance of early antiviral therapy.

Facial Palsy in Brainstem Lesions

Lesions involving the pons can affect the facial nerve nucleus or its intrapontine fibers. Such cases often present with **combined cranial nerve deficits**, long tract signs, or cerebellar features.

Examples include:

- Lateral pontine syndrome
- Demyelinating disorders such as multiple sclerosis
- Brainstem tumors

Recognition of these patterns is crucial for accurate neurological localization.

Synkinesis and Aberrant Regeneration



Synkinesis refers to involuntary facial movements occurring during voluntary actions, resulting from aberrant nerve regeneration. Neurologically, this occurs when regenerating axons innervate incorrect muscle groups.

Common manifestations include:

- Eye closure during smiling
- Mouth movement during blinking

Synkinesis is a marker of incomplete or misdirected neural recovery and often requires specialized rehabilitation.

Neuroplasticity and Recovery

Recovery from facial palsy depends on the degree of axonal injury and the capacity for neuroplasticity. Mild demyelinating lesions recover rapidly, while axonal degeneration leads to prolonged recovery and residual deficits.

Neurorehabilitation strategies aim to harness cortical reorganization and peripheral nerve regeneration to restore coordinated facial movement.

Prognostic Neurological Indicators

Neurological assessment plays a vital role in prognostication. Poor prognostic indicators include:

- Complete paralysis at onset

- Severe nerve degeneration on electrodiagnostic studies
- Delayed onset of recovery
- Presence of synkinesis

Conversely, early partial recovery suggests favorable neural regeneration.

Neurological Approach to Diagnosis

A comprehensive neurological evaluation of facial palsy includes:

- Detailed cranial nerve examination
- Assessment for central neurological signs
- Neuroimaging when indicated
- Correlation with electrodiagnostic findings

This systematic approach ensures accurate diagnosis and timely intervention.

Chapter Summary

Facial palsy is a neurological condition with diverse etiologies and presentations. Differentiation between central and peripheral causes is fundamental to clinical practice. Understanding the neurological mechanisms underlying facial nerve dysfunction enables accurate localization, effective management, and improved patient outcomes.

Further Reading:

Valls-Solé J. Facial nerve palsy and hemifacial spasm. Handb Clin Neurol. 2013;115:367-80. doi: 10.1016/B978-0-444-52902-2.00020-5. PMID: 23931790.

Estakhr M, Tabrizi R, Ghotbi Z, Shahabi S, Habibzadeh A, Bashi A, Borhani-Haghighi A. Is facial nerve palsy an early manifestation of COVID-19? A literature review. Am J Med Sci. 2022 Sep;364(3):264-273. doi: 10.1016/j.amjms.2022.04.010. Epub 2022 Apr 14. PMID: 35429449; PMCID: PMC9007824.

Lima MA, Silva MTT, Soares CN, Coutinho R, Oliveira HS, Afonso L, Espíndola O, Leite AC, Araujo A. Peripheral facial nerve palsy

associated with COVID-19. J Neurovirol. 2020 Dec;26(6):941-944. doi: 10.1007/s13365-020-00912-6. Epub 2020 Oct 2. PMID: 33006717; PMCID: PMC7531061.

Molaverdi G, Kamal Z, Safavi M, Shafiee A, Mozhgani SH, Ghobadi MZ, Goudarzvand M. Neurological complications after COVID-19: A narrative review. eNeurologicalSci. 2023 Dec;33:100485.

CHAPTER 4

Clinical Evaluation and Grading Systems

Dr. Darshpreet Kaur (PT)

Introduction

Clinical evaluation forms the cornerstone of diagnosis, prognosis, and management in facial palsy. In Bell's palsy, where the condition is largely idiopathic and self-limiting, a meticulous and timely clinical assessment is essential to determine severity, exclude alternative diagnoses, initiate appropriate treatment, and prevent long-term complications. This chapter outlines a systematic approach to clinical evaluation and discusses the commonly used grading systems for assessing facial nerve dysfunction.

Key Components of Clinical Evaluation

A comprehensive clinical evaluation of Bell's palsy begins with a detailed **patient history**, focusing on the onset, progression, and duration of facial weakness. Sudden onset with rapid progression over 24–48 hours is characteristic of Bell's palsy.

This is followed by careful **observation of facial symmetry** at rest and during voluntary movements. Particular attention should be paid to eyebrow elevation, forehead wrinkling, eye closure, smiling, and puckering of lips. Examination of the **nasolabial fold** and angle of the mouth provides valuable insight into lower facial muscle involvement.

A thorough **cranial nerve examination**, especially of the facial nerve (cranial nerve VII), is mandatory. Associated symptoms such as ear pain, hyperacusis, altered taste sensation, excessive tearing, or dry eye should be actively enquired about, as these help localize the lesion along the course of the facial nerve.

Importance of Early Clinical Evaluation

Early clinical evaluation is critical in Bell's palsy for several reasons. It helps determine the initial severity of nerve dysfunction, identify patients at risk of poor recovery, and guide early therapeutic decisions. More importantly, early assessment allows clinicians to **rule out other causes of facial paralysis**, such as stroke, tumors, infections, or systemic diseases.

Prompt diagnosis and grading also facilitate early initiation of pharmacological therapy and rehabilitation, which are key to preventing long-term sequelae such as facial contractures and synkinesis.

Differential Diagnosis in Facial Palsy

During clinical assessment, several differential diagnoses must be carefully excluded. These include:

- Central facial palsy due to cerebrovascular accidents
- Ramsay Hunt syndrome
- Facial nerve tumors
- Lyme disease
- Otitis media or mastoiditis

The presence of forehead sparing, additional neurological deficits, vesicular eruptions, or systemic symptoms should raise suspicion of alternative diagnoses and warrant further investigation.

Clinical Signs of Bell's Palsy

Typical clinical features of Bell's palsy include **unilateral facial droop**, inability to close the eye (lagophthalmos), flattening of the nasolabial fold, drooping of the corner of the mouth, and loss of forehead wrinkles on the affected side.

Bell's phenomenon, characterized by upward and outward movement of the eyeball during attempted eyelid closure, is commonly observed and has important clinical implications for eye care.

Role of Eye Examination

Eye examination is a vital component of facial palsy evaluation. Assessment of eyelid closure, corneal reflex, conjunctival dryness, and signs of irritation or exposure keratitis is essential. Inadequate eye protection can lead to corneal ulceration and permanent visual impairment.

Early identification of ocular involvement enables timely initiation of protective strategies such as lubrication, taping, or protective eyewear.

House–Brackmann Grading System

The **House–Brackmann (H–B) Grading System** is the most widely used clinical tool for grading facial nerve paralysis. It classifies facial function into six grades:

- **Grade I:** Normal facial function
- **Grade II:** Mild dysfunction
- **Grade III:** Moderate dysfunction
- **Grade IV:** Moderately severe dysfunction
- **Grade V:** Severe dysfunction
- **Grade VI:** Total paralysis

The simplicity and universal acceptance of this scale make it useful for initial assessment and communication among clinicians.

Table. House-Brackmann Facial Nerve Grading System

	Grade I <i>(Normal)</i>	Grade II <i>(Mild dysfunction)</i>	Grade III <i>(Mild- moderate dysfunction)</i>	Grade IV <i>(Moderate dysfunction)</i>	Grade V <i>(Severe dysfunction)</i>	Grade VI <i>(Complete paralysis)</i>
Resting Findings	Symmetric	Grossly symmetric	Grossly symmetric	Grossly symmetric	Grossly asymmetric (ptosis of the brow, lower eyelid, oral commissure, effacement of nasolabial fold)	Grossly asymmetric (often with a noticeable droop of the brow or oral commissure and effaced nasolabial fold, particularly in older patients)
Dynamic Findings	Symmetric	Mild asymmetry Full eye closure with gentle effort	Mild-moderate asymmetry Full eye closure only with full effort	Moderate brow elevation and smile asymmetry Incomplete eye closure	Severe brow elevation and smile asymmetry Incomplete eye closure	No movement Incomplete eye closure

From: [Bell Palsy](#)



StatPearls [Internet].
Treasure Island (FL): [StatPearls Publishing](#); 2025 Jan-.

[Copyright](#) © 2025, StatPearls Publishing LLC.

Limitations of the House–Brackmann System

Despite its widespread use, the House–Brackmann system has certain limitations. It is relatively subjective and lacks sensitivity to regional variations in facial movement, such as differences between eye and mouth function. It is also less effective in detecting subtle clinical changes during recovery.

Sunnybrook Facial Grading System

Sunnybrook Facial Grading System										
Resting Symmetry		Symmetry of Voluntary Movement					Synkinesis			
Compared to normal side		Degree of muscle EXCURSION compared to normal side					Rate the degree of INVOLUNTARY MUSCLE CONTRACTION associated with each expression			
Eye (choose one only)		Standard Expressions Forehead Wrinkle (FRD) Gentle eye closure (OCS) Open mouth smile (ZYG/RIS) Snarl (LLA/LLS) Lip Pucker (ODS/ODI) Severe Asymmetry Severe Asymmetry Moderate Asymmetry Mild Asymmetry Normal Symmetry Total					NONE: No synkinesis or slight movement MILD: Slight synkinesis MODERATE: Obvious but not debilitating synkinesis SEVERE: Debilitating synkinesis; Gross motor movement of facial muscles			
Cheek (naso-labial fold)										
normal	0	1	2	3	4	5	0	1	2	3
absent	2									
less pronounced	1									
more pronounced	1									
Mouth										
normal	0									
corner drooped	1									
corner pulled up/out	1									
Total										
Resting symmetry score		Total X 5								
Patient's name										
Dr.										
Date										
Vol mov't score		Resting symmetry score		Synk score		Composite score				

The Sunnybrook Facial Grading System (SBFGS) offers a more detailed and quantitative assessment of facial nerve function. It evaluates three key components:

- Resting facial symmetry
- Voluntary facial movements
- Synkinesis

The system generates a composite score out of 100, allowing precise documentation of facial function and recovery.

Clinical Utility of the Sunnybrook System

The Sunnybrook system is considered superior in many clinical scenarios due to its ability to identify regional differences, quantify improvement over time, and account for synkinesis. This makes it particularly useful in rehabilitation settings where tracking subtle changes is critical.

Combined Use of Grading Systems

Using both the House–Brackmann and Sunnybrook grading systems in clinical practice provides a comprehensive evaluation. While the House–Brackmann scale offers a broad overview of facial function, the Sunnybrook system delivers detailed insight into specific muscle groups and movement patterns. Together, they enhance clinical decision-making and documentation.

Frequency of Grading During Rehabilitation

Grading should ideally be repeated **every 2–3 weeks**, especially during the first 6–8 weeks following onset. Regular reassessment helps monitor progress, guide modifications in rehabilitation protocols, and objectively document recovery trends.

Prognostic Value of Grading Systems

Clinical grading systems also serve as valuable prognostic tools. Patients presenting with lower House–Brackmann grades (I–III) and higher baseline Sunnybrook scores generally experience faster and more complete recovery. Conversely, severe initial grades may indicate prolonged recovery and higher risk of residual deficits.

Integration with Electrodiagnostic Investigations

Clinical grading assesses visible facial performance, while electrodiagnostic studies such as electroneuronography and nerve conduction tests evaluate underlying nerve integrity. When used together, these tools provide a **comprehensive, full-spectrum assessment** of facial nerve function, enhancing accuracy in prognosis and treatment planning.

Chapter Summary

Clinical evaluation and grading systems play a pivotal role in the management of Bell’s palsy. A structured assessment, combined with appropriate grading tools, enables accurate diagnosis, effective monitoring, and informed clinical decision-making. Integration of

clinical findings with electrodiagnostic data ensures optimal patient outcomes.

Further Reading

Escalante DA, Malka RE, Wilson AG, Nygren ZS, Radcliffe KA, Ruhl DS, Vincent AG, Hohman MH. Determining the Prognosis of Bell's Palsy Based on Severity at Presentation and Electroneuronography. Otolaryngol Head Neck Surg. 2022 Jan;166(1):151-157.

Yilmaz OF, Oguz H. Ocular surface lesions in clinical grades of Bell's phenomenon. Med Hypothesis Discov Innov Ophthalmol. 2023 Winter;12(4):177-186

House JW, Brackmann DE. Facial nerve grading system. Otolaryngol Head Neck Surg. 1985 Apr;93(2):146-7.

Santosa KB, Fattah A, Gavilán J, Hadlock TA, Snyder-Warwick AK. Photographic Standards for Patients With Facial Palsy and Recommendations by Members of the Sir Charles Bell Society. JAMA Facial Plast Surg. 2017 Jul 01;19(4):275-281.

CHAPTER 5

Radiological Investigations in Facial Palsy

Dr. Vihag Raman

Introduction

Radiological investigations play a supportive but increasingly important role in the evaluation and management of facial palsy. Although Bell's palsy is primarily a clinical diagnosis, imaging is essential for excluding alternative etiologies, identifying the level of facial nerve involvement, assessing structural changes, and contributing to prognostic evaluation.

With advances in imaging technology, radiology has evolved from a purely exclusionary tool to an **active contributor to prognostication and rehabilitation planning**. This chapter reviews the role of ultrasonography, magnetic resonance imaging (MRI), high-resolution computed tomography (HRCT), and integrative protocols such as CESS and MRI–CESS in facial palsy.

Indications for Imaging in Facial Palsy

Imaging is not mandatory in every case of facial palsy. However, it is indicated in the following clinical situations:

- Atypical or progressive facial weakness
- Recurrent or bilateral facial palsy
- Delayed or incomplete recovery
- Presence of additional cranial nerve involvement
- Severe pain, trauma, or suspected infection
- Suspicion of central nervous system pathology
- Preoperative planning for surgical intervention

The choice of imaging modality depends on the suspected site and nature of pathology.

Ultrasonography of the Facial Nerve

Role and Technique

High-resolution ultrasonography allows direct visualization of the **extracranial segment of the facial nerve**, particularly the main trunk as it emerges from the stylomastoid foramen and courses through the parotid gland. Using high-frequency linear probes, the facial nerve can be identified and measured in real time.

Findings in Bell's Palsy

In Bell's palsy, ultrasonography commonly demonstrates:

- Increased facial nerve diameter
- Altered echotexture indicating edema
- Asymmetry compared to the contralateral side

These findings reflect inflammatory swelling extending distally from the site of nerve injury.

Prognostic Significance

Ultrasound has emerged as a valuable prognostic tool. Measurement of the distal facial nerve diameter has shown:

- **Positive predictive value of 100%** for favorable recovery
- **Negative predictive value of approximately 77%** for poor recovery at three months

Some studies have demonstrated ultrasound to be superior to conventional electrodiagnostic tests in early recovery prediction.

Magnetic Resonance Imaging (MRI)

Role of MRI

MRI is the modality of choice for evaluating the **intracranial and intratemporal segments** of the facial nerve. It is particularly useful for identifying central causes of facial palsy and excluding serious pathologies.

MRI Findings in Bell's Palsy

Typical MRI features include:

- Contrast enhancement of the facial nerve
- Predominant involvement of the **labyrinthine segment** and **geniculate ganglion**
- Absence of mass lesions, infarction, or demyelination

Contrast-enhanced MRI helps differentiate Bell's palsy from tumors, stroke, and inflammatory disorders.

High-Resolution Computed Tomography (HRCT)

Role of HRCT Temporal Bone

HRCT is primarily used to evaluate **bony anatomy** of the facial canal and middle ear structures. It is indicated in:

- Facial palsy following trauma
- Chronic otitis media or mastoid disease
- Congenital anomalies
- Preoperative assessment for facial nerve decompression

In Bell's palsy, HRCT is not routinely required but may reveal anatomical predispositions such as a narrow facial canal.

Imaging in Ramsay Hunt Syndrome and Other Conditions

MRI plays a crucial role in diagnosing Ramsay Hunt syndrome, often demonstrating:

- Facial nerve and geniculate ganglion enhancement
- Involvement of adjacent cranial nerves
- Inner ear inflammation

Imaging is also essential in facial palsy caused by tumors, infections, or brainstem lesions, where it guides definitive management.

MRI–CESS Protocol

The **MRI–CESS protocol** expands the CESS framework by incorporating MRI findings.

Components

- MRI assessment of proximal facial nerve inflammation
- Ultrasound evaluation of distal nerve edema
- Electrical stimulation response of facial muscles

Clinical Value

This integrated approach allows:

- Multilevel localization of nerve pathology
- Early prognostic stratification
- Informed rehabilitation planning

Marked MRI enhancement with poor stimulation response suggests severe nerve injury, whereas mild enhancement with preserved response indicates favorable recovery.

Clinical Integration of Imaging Modalities

Each imaging modality contributes uniquely:

- **Ultrasound** – Extracranial nerve assessment and early prognosis
- **MRI** – Central and intratemporal pathology
- **HRCT** – Bony anatomy and surgical planning
- **CESS / MRI–CESS** – Integrated prognostic and rehabilitation guidance

Judicious integration of these tools enhances diagnostic accuracy and individualized care.

Limitations of Imaging

- Operator dependency (ultrasound)
- Cost and accessibility (MRI)

- Limited sensitivity in mild cases

Imaging findings must always be interpreted in conjunction with clinical evaluation.

Physiotherapist-Oriented Summary

What Physiotherapists Need to Know

- Imaging supports **rehabilitation decision-making**, not diagnosis
- **Ultrasound** is the most relevant imaging tool for physiotherapists
- Increased facial nerve diameter indicates inflammation and possible delayed recovery
- MRI helps explain atypical or slow recovery patterns
- HRCT is mainly relevant in post-traumatic or post-surgical cases

How Imaging Guides Rehabilitation

- Mild nerve swelling + good muscle response → early active therapy
- Moderate findings → graded, closely monitored rehabilitation
- Severe imaging abnormalities → protective phase, delayed strengthening

Key Takeaway

For physiotherapists, imaging findings should be used to:

- Set realistic recovery timelines
- Prevent over-aggressive or delayed therapy
- Individualize facial rehabilitation protocols
- Improve patient education and compliance

When integrated with clinical grading systems and electrodiagnostic data, radiological investigations enhance **precision, safety, and outcomes in facial palsy rehabilitation**.

Further reading:

Seok JI, Park JH, Park JA, Do Y. Contrast-enhanced MRI findings of patients with acute Bell palsy within 7 days of symptom onset: A retrospective study. *Medicine (Baltimore)*. 2023 Dec 01;102(48):e36337.

Lo YL, Fook-Chong S, Leoh TH, Dan YF, Lee MP, Gan HY, Chan LL. High-resolution ultrasound in the evaluation and prognosis of Bell's palsy. *Eur J Neurol*. 2010 Jun 01;17(6):885-9.

Celik O, Eskiizmir G, Pabescu Y, Ulkumen B, Toker GT. The role of facial canal diameter in the pathogenesis and grade of Bell's palsy: a study by high resolution computed tomography. *Braz J Otorhinolaryngol*. 2017 May-Jun;83(3):261-268.

Gupta S, Mends F, Hagiwara M, Fatterpekar G, Roehm PC. Imaging the facial nerve: a contemporary review. *Radiol Res Pract*. 2013;2013:248039.

CHAPTER 6

Electrophysiological Tests Used in Facial Palsy


Dr. (Prof) Gunjan Kumar
Dr. Darshpreet Kaur (PT)

Electroneurography (ENoG)

Purpose and Clinical Role

Electroneurography (ENoG) is the **most reliable electrophysiological test for prognostication** in acute complete facial paralysis. It quantifies the degree of facial nerve degeneration by measuring the compound muscle action potential (CMAP) generated by electrical stimulation of the facial nerve.

Originally described by Esslen and later popularized by Fisch, ENoG was developed to improve prognostic accuracy in facial palsy. The term *Electroneurography (ENoG)* is now recommended for routine clinical use.



Methodology

- Supramaximal electrical stimulation is delivered to the facial nerve at the **stylomastoid foramen**
- CMAPs are recorded from facial muscles such as the nasolabial fold or orbicularis oris
- The CMAP amplitude on the affected side is compared with the unaffected side, which serves as an internal control
- Recording electrodes placed on each side of the face. Stimulator placed in front of the ear.

CMAP amplitude reflects the number of **viable, synchronously firing axons**. A reduction in amplitude indicates Wallerian degeneration.

Interpretation

- 90% degeneration within 10–14 days → **poor prognosis**
- Preservation of CMAP indicates neurapraxia with potential for recovery
- Absence of transmission suggests axonotmesis or neurotmesis

Optimal Timing

ENoG is most reliable when performed between **Day 3 and Day 21** post-onset, with peak prognostic value between **Day 7 and Day 14**.

Electromyography (EMG)

Purpose

Electromyography evaluates **voluntary motor unit recruitment** and signs of **reinnervation**, making it particularly useful in the subacute and recovery phases of Bell's palsy.

Methodology

- Needle EMG electrodes are inserted into selected facial muscles
- Motor unit potentials (MUPs) are analyzed during voluntary contraction

Key Findings

- Absence of voluntary activity after Day 10–14 → **poor prognosis**
- Presence of polyphasic reinnervation potentials → **ongoing recovery**

Timing

EMG is most informative **after 10–14 days**, when denervation and reinnervation changes become evident.


Blink Reflex Test

Purpose

The blink reflex evaluates the **trigemino-facial reflex arc**, involving:

- Trigeminal nerve (CN V)
- Facial nerve (CN VII)
- Brainstem interneuronal connections

It is especially useful for **localizing lesions** and differentiating **central versus peripheral facial palsy**.



Methodology

- Electrical stimulation of the supraorbital nerve (V1)
- Bilateral recording from orbicularis oculi muscles

Responses Recorded

- **R1:** Early ipsilateral response (~10 ms; pontine circuit)
- **R2i:** Late ipsilateral response (~30 ms)
- **R2c:** Late contralateral response (~30–40 ms)

Findings in Bell's Palsy

- Absent or delayed R1 and R2 responses on the affected side
- Normal blink reflex helps exclude central causes such as stroke or multiple sclerosis

Nerve Excitability Test (NET) and Maximal Stimulation Test (MST)

These are **early bedside screening tests** now largely superseded by ENoG.

- **NET:** Compares minimum current needed to elicit visible contraction on both sides
- **MST:** Compares symmetry of muscle contraction at maximal stimulation

A side-to-side difference >3.5 mA suggests degeneration, but these tests are subjective and have limited prognostic accuracy.

Comparative Overview of Electrodiagnostic Tests

Test	Best Time	Primary Use	Prognostic Value
ENoG	Day 3–21	Quantify degeneration	High
EMG	After Day 10–14	Reinnervation assessment	High
Blink Reflex	Anytime	Lesion localization	Moderate
NET / MST	Day 1–5	Bedside screening	Low–Moderate

Degeneration Index (DI)

Definition and Clinical Importance

The **Degeneration Index (DI)** is a quantitative measure derived from ENoG that represents the **percentage loss of facial nerve function** on the affected side compared with the unaffected side. It is a critical prognostic indicator and assists in identifying patients with severe nerve injury.

Formula

$$DI\ (\%) = \frac{1 - \text{CMAP amplitude (affected side)}}{\text{CMAP amplitude (unaffected side)}} \times 100$$

Example:

Unaffected side = 3.5 mV

Affected side = 0.3 mV

DI = 91.4% → Severe degeneration, poor prognosis

Temporal Evolution of DI in Bell’s Palsy

Time Post-Onset	Pathophysiology	DI Trend	Interpretation
0–48 hours	No Wallerian degeneration	Low DI	Damage underestimated
Day 3–5	Degeneration begins	Rising DI	Early prognostic phase
Day 5–10	Peak degeneration	Rapid DI rise	Critical window
Day 10–14	Maximal degeneration	DI plateaus	Best prognostic value
>21 days	Reinnervation begins	DI unreliable	Prognosis confounded

Grading of Degeneration Index

DI Value	Severity	Prognosis
<50%	Mild	Good
50–90%	Moderate	Variable
>90%	Severe	Poor

Clinical Decision-Making Based on DI

- **Corticosteroids** should be initiated regardless of DI
- Low DI (<50%) → excellent response to medical therapy
- High DI (>90%) → indicates severe axonal injury
- **Surgical decompression** is rarely indicated and is reserved for extreme cases
- Current guidelines do **not recommend routine surgery**, even with high DI

Timing: The Most Critical Factor in DI Interpretation

Accurate prognostication depends on **correct timing** of testing.

Timing	Reliability	Reason
<72 hours	✗ Poor	Degeneration incomplete
Day 3–10	✓ Good	Active degeneration
Day 10–14	✓✓ Best	Peak axonal loss
>21 days	⚠ Reduced	Regeneration interferes

Testing outside the optimal window may lead to false reassurance or misinterpretation.

Guideline Recommendations

Organization	Recommendation
AAO-HNSF (2020)	Use ENoG in complete palsy within 3–21 days
AAN	Supports ENoG for assessing axonal degeneration
EFNS	Recommends ENoG when surgery is considered
NICE (UK)	Emphasizes clinical management; no routine ENoG

Chapter Summary

Electrodiagnostic investigations provide objective, time-sensitive insights into facial nerve injury in Bell’s palsy. ENoG remains the gold standard for prognostication in complete paralysis, while EMG and blink reflex testing complement localization and recovery assessment. Correct timing and integration with clinical findings are essential for accurate interpretation.

Further Reading:

May M, Schaitkin BM. *The Facial Nerve*. New York, NY: Thieme; 2000.

Pauna, H. F., Silva, V. A. R., Lavinsky, J., Hyppolito, M. A., Vianna, M. F., Gouveia, M. C. L., Monsanto, R. C., Polanski, J. F., Silva, M. N. L., Soares, V. Y. R., Sampaio, A. L. L., Zanini, R. V. R., Abrahão, N. M., Guimarães, G. C., Chone, C. T., & Castilho, A. M. (2023). Task force of the Brazilian Society of Otology—Evaluation and management of peripheral facial palsy. *Brazilian Journal of Otorhinolaryngology*, 89(4), 459–475.

Baugh, R. F., Basura, G. J., Ishii, L. E., Schwartz, S. R., Drumheller, C. M., Burkholder, R., ... & Rosenfeld, R. M. (2020). Clinical practice guideline: Bell's palsy (update). *Otolaryngology–Head and Neck Surgery*, 162(2_suppl), S1–S45.

American Academy of Audiology. *Audiology: Scope of Practice Position Statements*. See <https://www.audiology.org/Pages/default.aspx> (online Dec. 2000).

Kimura, M., Nakagawa, I., Miimai, H., Hamada, H., Sakai, A., and Yasuuiji, M.: Evaluation with blink reflex of bilateral facial palsy. *Masui*, 49:2, 159-162. Feb. 2000.

Fisch U. Prognostic value of electrical tests in acute facial paralysis. *Am J Otol*. 1984 Oct;5(6):494-8.

Guntinas-Lichius O, Volk GF, Olsen KD, Mäkitie AA, Silver CE, Zafereo ME, Rinaldo A, Randolph GW, Simo R, Shaha AR, Vander Poorten V, Ferlito A. Facial nerve electrodiagnostics for patients with facial palsy: a clinical practice guideline. *Eur Arch Otorhinolaryngol*. 2020 Jul;277(7):1855-1874.

CHAPTER 7

Pharmacological and Acute Phase Management of Facial Palsy

Dr. (Prof) Ajay Kumar Singh

Dr. (Prof) Gunjan Kumar

Introduction

The acute phase of facial palsy, particularly Bell's palsy, represents a critical therapeutic window during which timely pharmacological intervention and supportive care can significantly influence recovery outcomes. Early management aims to reduce facial nerve inflammation, prevent irreversible axonal damage, protect ocular structures, and provide psychological reassurance to the patient. This chapter outlines evidence-based pharmacological strategies and essential acute phase care principles.

Corticosteroid Therapy

Rationale

Inflammation and edema of the facial nerve within the narrow facial canal are central to the pathophysiology of Bell's palsy. Corticosteroids reduce nerve swelling, improve microcirculation, and limit secondary ischemic injury, thereby enhancing the likelihood of functional recovery.

Indications

- All patients with acute idiopathic peripheral facial palsy
- Most effective when initiated within 72 hours of symptom onset
- Beneficial across mild, moderate, and severe presentations

Commonly Used Regimens

Typical adult regimens include:

- **Prednisolone 1 mg/kg/day** (maximum 60 mg/day) for 5–7 days
- Followed by gradual tapering over the next 5–7 days

Alternative equivalent corticosteroids may be used based on clinical judgment and patient tolerance.

Clinical Evidence

Multiple randomized controlled trials and meta-analyses have demonstrated that early corticosteroid therapy:

- Significantly improves complete recovery rates
- Reduces long-term complications such as synkinesis and residual weakness

Corticosteroids are currently considered the **cornerstone of acute Bell’s palsy management**.

Precautions and Contraindications

- Diabetes mellitus (monitor blood glucose closely)
- Hypertension
- Peptic ulcer disease
- Active infections

Risk–benefit assessment and patient-specific modification are essential.

Antiviral Medications

Rationale

Viral reactivation—particularly herpes simplex virus type 1—has been implicated in the etiology of Bell’s palsy. Antiviral agents aim to suppress viral replication and potentially reduce nerve inflammation.

Indications

- Moderate to severe facial palsy (House–Brackmann Grade IV–VI)
- Suspected **Ramsay Hunt syndrome**
- Presence of severe pain or vesicular eruptions
- As an adjunct to corticosteroids (not as monotherapy)

Commonly Used Antivirals

- **Acyclovir**: 400 mg five times daily for 7–10 days
- **Valacyclovir**: 1 g three times daily for 7 days

Valacyclovir is often preferred due to better bioavailability and dosing convenience.

Clinical Evidence

Current evidence suggests that:

- Antivirals alone are **not effective**
- Combination therapy with corticosteroids may offer **modest benefit**, particularly in severe cases
- Corticosteroids remain essential regardless of antiviral use

Eye Care and Corneal Protection

Importance of Eye Protection

In Bell's palsy, weakness of the orbicularis oculi muscle leads to **incomplete eyelid closure (lagophthalmos)** and reduced blink reflex, increasing the risk of corneal dryness, exposure keratitis, and ulceration.

Assessment

Eye evaluation should include:

- Degree of eyelid closure
- Corneal reflex
- Signs of dryness, redness, or irritation

Eye Care Measures



- **Lubricating eye drops** (artificial tears) during the day
- **Lubricating ointment** at night
- **Eye patching or taping** during sleep
- Protective glasses in outdoor environments

Early ophthalmology referral is indicated if corneal involvement is suspected.

Clinical Significance

Adequate eye care is a **preventive emergency**. Neglect can result in permanent visual impairment, even when facial nerve recovery is otherwise satisfactory.

Acute Phase Patient Counseling

Purpose of Counseling

The sudden onset of facial paralysis can be distressing for patients. Clear communication during the acute phase helps:

- Reduce anxiety
- Improve treatment compliance
- Set realistic expectations

Key Counselling Points

Patients should be informed that:

- Bell's palsy is **usually temporary**
- Most patients show improvement within **3–6 weeks**
- Complete recovery is common, especially with early treatment
- Facial exercises are **not recommended aggressively** in the very acute phase unless guided by a professional
- Eye protection is crucial and must be followed strictly

Lifestyle and Activity Advice

- Avoid facial overuse or forceful movements initially
- Maintain adequate nutrition and hydration
- Adhere strictly to medication schedules
- Attend scheduled follow-ups for reassessment

Psychological Support

Reassurance and emotional support are essential components of care. Addressing cosmetic concerns, social anxiety, and fear of permanent damage improves overall outcomes and patient satisfaction.

Integration with Rehabilitation

Pharmacological and acute phase management sets the foundation for successful rehabilitation. Early reduction of inflammation and prevention of complications allow physiotherapy and neuromuscular re-education to be introduced at the appropriate time with better functional results.

Chapter Summary

Early pharmacological intervention—particularly corticosteroid therapy—combined with appropriate antiviral use, meticulous eye care, and effective patient counseling forms the cornerstone of acute Bell's palsy management. Timely and holistic acute phase care not only

improves recovery rates but also prevents long-term complications and optimizes rehabilitation outcomes.

Further Reading:

Hato N, Sawai N, Teraoka M, Wakisaka H, Takahashi H, Hinohira Y, Gyo K. Valacyclovir for the treatment of Bell's palsy. Expert Opin Pharmacother. 2008 Oct;9(14):2531-6.

Sullivan FM, Swan IR, Donnan PT, Morrison JM, Smith BH, McKinstry B, Davenport RJ, Vale LD, Clarkson JE, Hammersley V, Hayavi S, McAteer A, Stewart K, Daly F. Early treatment with prednisolone or acyclovir in Bell's palsy. N Engl J Med. 2007 Oct 18;357(16):1598-607.

Hato N, Yamada H, Kohno H, Matsumoto S, Honda N, Gyo K, Fukuda S, Furuta Y, Ohtani F, Aizawa H, Aoyagi M, Inamura H, Nakashima T, Nakata S, Murakami S, Kiguchi J, Yamano K, Takeda T, Hamada M, Yamakawa K. Valacyclovir and prednisolone treatment for Bell's palsy: a multicenter, randomized, placebo-controlled study. Otol Neurotol. 2007 Apr;28(3):408-13.

CHAPTER 8


Plastic and Reconstructive Surgical Approaches in Facial Palsy

Dr. Sarsij Sharma

Introduction

Surgical management of facial palsy is reserved for selected cases where spontaneous recovery is unlikely or incomplete, or where long-standing paralysis has resulted in functional and aesthetic deficits. Successful facial nerve surgery requires an in-depth understanding of **facial nerve anatomy**, careful patient selection, and appropriate timing of intervention. This chapter reviews surgically relevant anatomy, facial nerve landmarks, reconstructive options, and decision-making principles in facial reanimation.

Surgically Relevant Anatomy of the Facial Nerve



Source: Lam A Q, Tran Phan Chung T, Tran Viet L, et al. (March 03, 2022) The Anatomic Landmark Approach to Extratemporal Facial Nerve Repair in Facial Trauma. Cureus 14(3): e22787. doi:10.7759/cureus.22787

Key Anatomical Landmarks for Identifying the Facial Nerve Trunk

Accurate identification of the facial nerve trunk is fundamental during parotidectomy and facial nerve exploration. Several reliable anatomical landmarks are used:

1. **Tragal Pointer**

The deep end of the tragal cartilage forms a pointed structure known as the tragal pointer. The facial nerve lies approximately **1 cm deep and inferior** to this point. An avascular plane anterior to the tragus allows safe dissection toward this landmark.

2. **Posterior Belly of Digastric Muscle**

The facial nerve trunk emerges just **superior to the upper border** of the posterior belly of the digastric muscle, making it a consistent and dependable landmark during surgery.

3. **Tympanomastoid Suture**

Tracing the anterior border of the mastoid process superiorly reveals its junction with the tympanic bone. The facial nerve lies at the **bisector of the angle** formed at this junction.

4. **Styloid Process**

Palpation of the styloid process between the mastoid bone and posterior border of the mandible helps identify the nerve, which lies **just lateral** to the process.

Pes Anserinus

The **pes anserinus** refers to the point where the facial nerve trunk divides into its major branches within the parotid gland. Identification of this branching point is crucial for preserving nerve integrity during parotid and reconstructive procedures.

Facial Nerve “Danger Zones” in Facial Surgery

During SMAS elevation in facelift procedures, the facial nerve traverses predictable planes but crosses fascial layers in specific regions known as **danger zones**, where nerve injury risk is high:

- **Zone 1:** Greater auricular nerve emerging from beneath the sternocleidomastoid muscle
- **Zone 2:** Temporal branch running beneath the temporoparietal fascia–SMAS layer
- **Zone 3:** Marginal mandibular branch where the platysma–SMAS layer thins
- **Zone 4:** Zygomatic and buccal branches anterior to the parotid gland
- **Zone 5:** Supraorbital and supratrochlear nerves at their foramina
- **Zone 6:** Infraorbital nerve at the infraorbital foramen
- **Zone 7:** Mental nerve at its exit below the second mandibular premolar

Dissection in these zones should be either avoided or performed with meticulous care.

Vulnerability of the Temporal Branch

The **temporal (frontal) branch** is the most vulnerable during temporal or frontal surgical approaches. It runs on the undersurface of the superficial temporal fascia. Safe dissection is performed either superficial or deep to the deep temporal fascia to avoid nerve injury.

Dynamic Facial Reanimation Techniques

Dynamic reanimation aims to restore **active facial movement**, particularly smiling.

Available Options

- **Nerve Transfers** (using contralateral facial nerve or other cranial nerves)
- **Cross-Facial Nerve Grafting (CFNG)**
- **Free Functional Muscle Transfer** (commonly gracilis muscle)
- **Regional Muscle Transfers** (e.g., temporalis muscle transfer)

Decision-Making: Nerve Grafting vs Muscle Transfer

The **duration of facial paralysis** is the principal determinant:

- **< 12 months:** Facial muscles are usually viable → nerve repair or grafting preferred
- **12–24 months:** Partial muscle viability → combined strategies may be required
- **> 24 months:** Muscles are irreversibly atrophic → muscle replacement indicated

Role of the Sural Nerve in Facial Nerve Repair

The **sural nerve** is commonly harvested as a **cable nerve graft** to bridge gaps between the proximal facial nerve stump and distal branches.

In **cross-facial nerve grafting**, the sural nerve serves as a conduit allowing axons from the contralateral healthy facial nerve to regenerate across the face and reinnervate paralyzed muscles over **4–8 months**.

Intraoperative Facial Nerve Assessment

Facial nerve function is assessed intraoperatively using **microbipolar electrical stimulation**. Zygomaticobuccal branches medial to the parotid gland are individually stimulated to confirm viability and select appropriate donor branches.

Branches producing a **pure smile without orbicularis oculi activation** are preferred during reanimation procedures.

Cross-Facial Nerve Grafting: Donor and Recipient Selection

The donor nerve is typically the **contralateral facial nerve**. Zygomaticobuccal branches producing smile are identified and selected. Usually **two to four branches** are used to optimize axonal input while preserving donor-side function.

Surgical Management of Bell's Palsy

Indications for Surgical Decompression

Surgical decompression is rarely indicated and considered only when:

- Medical therapy fails
- ENoG shows **no nerve activity after two weeks**
- Severe axonal degeneration is demonstrated within three weeks
- Significant nerve edema is present within the fallopian canal

Surgical Approaches

- **Transmastoid Approach:** For tympanic and mastoid segment involvement
- **Middle Cranial Fossa Approach:** For labyrinthine segment involvement
- **Combined Approach:** For complex temporal bone fractures

Decompression is typically performed over **180 degrees** of the nerve circumference.

Timing of Nerve Repair

Early nerve repair results in:

- Improved axonal regeneration
- Reduced fibrosis and scarring
- Lower risk of aberrant regeneration

Delayed repair is associated with inferior outcomes and higher rates of synkinesis.

Masseter Nerve Transfer and the Babysitter Procedure

The **masseter nerve** plays a crucial role in facial reanimation, particularly as part of the **babysitter procedure**. While cross-facial nerve grafts regenerate slowly, an ipsilateral motor nerve (masseter or hypoglossal) provides temporary innervation to preserve muscle tone.

In cases of bilateral facial paralysis or absent facial nerve donors, the masseter nerve serves as the primary motor source for muscle reinnervation.

Free Functional Muscle Transfer: Gracilis Muscle

Anatomical Considerations

- **Axis:** Two fingerbreadths anterior to adductor longus tendon
- **Origin:** Pubic tubercle
- **Insertion:** Tibial condyle
- **Vascular Pedicle:** Located ~10 cm below pubic tubercle
- **Nerve Supply:** Anterior branch of obturator nerve, entering obliquely at 45°

Gracilis muscle is preferred due to its reliable anatomy and minimal donor-site morbidity.

Chapter Summary

Plastic and reconstructive surgery offers powerful solutions for facial palsy when spontaneous recovery is incomplete or absent. Mastery of facial nerve anatomy, careful selection of reconstructive techniques, and appropriate timing are critical to successful outcomes. Dynamic reanimation strategies—particularly nerve grafting, muscle transfer, and nerve transfers—restore function, symmetry, and quality of life in patients with facial paralysis.

Further Reading:

Seckel BR. Facial Danger Zones: Avoiding Nerve Injury in Facial Plastic Surgery. St. Louis: Quality Medical; 1994.

Myckatyn TM, MacKinnon SE. A review of facial nerve anatomy. Semin Plast Surg. 2004;18:5–12.

Wilhemi BJ, Mowlavi A, Neumeister MW. The safe face lift with bony anatomic landmarks to elevate the SMAS. Plast Reconstr Surg. 2003;111:1723–1726.

Ishikawa Y. An anatomical study on the distribution of the temporal branch of the facial nerve. J Craniomaxillofac Surg. 1990;18:287–292.

Agarwal CA, Mendenhall SD 3rd, Foreman KB, et al. The course of the frontal branch of the facial nerve in relation to fascial planes: an anatomic study. Plast Reconstr Surg. 2010;125:532– 537.

Manktelow RT. Microvascular Reconstruction. Anatomy, Applications, and Surgical Technique. New York: Springer-Verlag; 1986.

CHAPTER 9

Physiotherapy and Rehabilitation Techniques in Facial Palsy

Dr. Darshpreet Kaur (PT)

Dr. Nidhi Billore (PT)

Introduction

Physiotherapy plays a central role in the functional recovery of patients with facial palsy. While pharmacological and surgical interventions address the underlying pathology, rehabilitation focuses on **restoring coordinated facial movement, preventing secondary complications, and improving psychosocial well-being**. A structured, stage-wise physiotherapy program is essential to optimize outcomes and minimize long-term sequelae such as synkinesis, contractures, and facial asymmetry.

This chapter outlines the principles, techniques, and progression of facial rehabilitation across different stages of recovery.

Principles of Facial Rehabilitation

Effective facial rehabilitation is based on the following core principles:

- Individualized, stage-specific intervention
- Emphasis on **quality of movement rather than force**
- Prevention of maladaptive motor patterns
- Integration of sensory feedback and motor control
- Continuous reassessment and graded progression

Early overuse or inappropriate exercises can worsen outcomes; hence, physiotherapy must be **timely, precise, and supervised**.

Phases of Rehabilitation in Facial Palsy

1. Acute Phase (0–2 Weeks)

Goals

- Protect ocular structures
- Reduce edema and discomfort
- Maintain soft tissue mobility
- Prevent learned non-use

Physiotherapy Interventions

- Patient education and reassurance
- Gentle facial massage to promote circulation
- Passive range of motion for facial muscles
- Assisted eye closure techniques
- Postural correction for cervical and cranial alignment

Note: Forceful facial exercises and electrical stimulation are avoided in the early acute phase.

2. Subacute Phase (2–6 Weeks)

Goals

- Facilitate voluntary muscle activation
- Improve facial symmetry
- Begin neuromuscular re-education

Physiotherapy Interventions

- Assisted-to-active facial movements
- Mirror-guided exercises
- Gentle motor facilitation techniques
- Proprioceptive neuromuscular facilitation (PNF) principles adapted for facial muscles
- Soft tissue mobilization

Progression is guided by clinical grading systems and neuromuscular responsiveness.

3. Recovery Phase (6 Weeks–6 Months)

Goals

- Enhance coordination and strength
- Normalize facial expressions
- Prevent synkinesis

Physiotherapy Interventions

- Targeted facial muscle exercises
- Selective activation training
- Inhibition of unwanted movements
- Functional expression training (smiling, blinking, speech-related movements)
- Biofeedback-assisted retraining

This phase requires meticulous supervision to ensure correct motor patterns.

4. Chronic Phase (>6 Months)

Goals

- Manage residual weakness or synkinesis
- Improve facial symmetry at rest and during movement
- Enhance quality of life

Physiotherapy Interventions

- Synkinesis control strategies
- Relaxation and stretching of hyperactive muscles
- Advanced neuromuscular retraining
- Integration with adjunctive medical or surgical treatments if required

Facial Muscle Exercises

Facial exercises aim to retrain coordinated movement rather than increase muscle strength.

Key Principles

- Slow, controlled movements
- Low repetitions with high precision
- Avoid mass movements
- Emphasize symmetry

Exercises are tailored to individual muscle groups such as orbicularis oculi, zygomaticus major, orbicularis oris, and frontalis.

Mirror Therapy and Visual Feedback



Mirror therapy enhances cortical reorganization by providing visual feedback of symmetrical facial movements. It helps:

- Improve motor awareness
- Reduce abnormal movement patterns
- Enhance voluntary control

Short, frequent sessions are preferred over prolonged practice.

Biofeedback Techniques



Biofeedback, including surface electromyography (sEMG), allows patients to visualize muscle activation and selectively recruit specific muscles. It is particularly effective in:

- Preventing synkinesis
- Refining movement quality
- Enhancing patient engagement

Low-Level Laser Therapy (LLLT)



Rationale

LLLT is used to:

- Reduce inflammation
- Improve microcirculation
- Facilitate nerve regeneration

Application

- Applied along the course of the facial nerve and affected muscles
- Dosage and parameters should follow standardized protocols

LLLT serves as an adjunct to active rehabilitation.

Electrical Stimulation

Role and Controversy

Electrical stimulation remains a debated modality in facial palsy. When used appropriately:

- It may help maintain muscle excitability
- Prevent disuse atrophy

Guidelines for Use

- Low intensity, short duration
- Avoid in early acute phase
- Avoid indiscriminate or high-intensity stimulation
- Always combined with voluntary effort

Improper use may increase synkinesis risk.

Acoustic Shock Wave Therapy



Acoustic shock wave therapy is an emerging modality that may:

- Improve tissue pliability
- Enhance neuromuscular responsiveness
- Reduce fibrosis in chronic cases

It is primarily used as an adjunct in selected patients with delayed recovery.

Soft Tissue Mobilization and Stretching



Manual techniques help:

- Reduce muscle stiffness
- Improve circulation
- Address secondary myofascial tightness

Stretching is particularly useful in managing chronic synkinesis and contractures.

Management of Synkinesis

Synkinesis results from aberrant nerve regeneration and requires specialized rehabilitation strategies.

Interventions

- Movement inhibition techniques
- Muscle relaxation training
- Selective motor control exercises
- Biofeedback-assisted retraining

Early identification and targeted therapy are critical.

Psychosocial Considerations

Facial palsy significantly impacts self-image and social interaction. Physiotherapists should:

- Provide empathetic counseling
- Encourage patient adherence
- Address fear of permanent disfigurement

Psychological reassurance enhances rehabilitation outcomes.

Integration with Imaging and Electrodiagnostics

Rehabilitation planning should be informed by:

- Clinical grading systems
- Electrodiagnostic findings
- Imaging and CESS/MRI–CESS protocols

This ensures appropriate timing, intensity, and progression of therapy.

Outcome Measures in Rehabilitation

Progress should be documented using:

- House–Brackmann scale
- Sunnybrook Facial Grading System
- Patient-reported outcome measures

Objective tracking supports evidence-based practice.

Chapter Summary

Physiotherapy is a cornerstone in the comprehensive management of facial palsy. A structured, phase-wise rehabilitation program emphasizing neuromuscular re-education, movement quality, and prevention of maladaptive patterns leads to optimal functional recovery. Integration of adjunctive modalities, patient education, and continuous reassessment ensures individualized, effective care and improved quality of life.

Further Reading:

Tuncay F, Borman P, Taşer B, Ünlü İ, Samim E. Role of electrical stimulation added to conventional therapy in patients with idiopathic facial (Bell) palsy. Am J Phys Med Rehabil. 2015 Mar;94(3):222-8.

Di Pietro A, Cameron M, Campana V, Leyes L, Zalazar Cinat JAI, Lochala C, Johnson CZ, Hildebrand A, Loyo M. Efficacy of adding selective electrical muscle stimulation to usual physical therapy for Bell's palsy: immediate and six-month outcomes. Eur J Transl Myol. 2023 Oct 24;33(4)

Cronin GW, Steenerson RL. The effectiveness of neuromuscular facial retraining combined with electromyography in facial paralysis rehabilitation. Otolaryngol Head Neck Surg. 2003 Apr;128(4):534-8.

Nakamura K, Toda N, Sakamaki K, Kashima K, Takeda N. Biofeedback rehabilitation for prevention of synkinesis after facial palsy. Otolaryngol Head Neck Surg. 2003 Apr;128(4):539-43.

Lindsay RW, Robinson M, Hadlock TA. Comprehensive facial rehabilitation improves function in people with facial paralysis: a 5-year experience at the Massachusetts Eye and Ear Infirmary. Phys Ther. 2010 Mar;90(3):391-7.

CHAPTER 10

Proprietary Rehabilitation Protocol and Clinical Data

Dr. Darshpreet Kaur (PT)

Dr. (Prof) Gunjan Kumar

Introduction

Despite advances in pharmacological and surgical management, variability in rehabilitation outcomes remains a major challenge in Bell's palsy. Differences in severity, timing of intervention, and inappropriate use of modalities—particularly electrical stimulation—often contribute to delayed recovery and long-term complications such as synkinesis and facial contractures.

To address these gaps, **Darsh–Gunjan's Bell's Palsy Treatment Protocol©** was developed as a structured, severity-based, multimodal rehabilitation framework. This protocol integrates **objective electrodiagnostic data** with **standardized physiotherapy techniques**, ensuring personalized yet reproducible care.

Darsh–Gunjan's Bell's Palsy Treatment Protocol©

Darsh–Gunjan's Bell's Palsy Treatment Protocol© is a **standardized yet individualized multimodal physiotherapy regimen**, formally registered with the **Government of India in May 2024**. The protocol is uniquely designed around **ENoG-based severity stratification** combined with validated clinical grading systems.

The central philosophy of the protocol is that **objective quantification of nerve degeneration must guide rehabilitation intensity and modality selection**, rather than relying solely on visual clinical assessment.



INTELLECTUAL PROPERTY INDIA
भारत सरकार, विज्ञान, सूचना एवं संचार विभाग
भारत सरकार, विज्ञान, सूचना एवं संचार विभाग



Extracts from the Register of Copyrights



प्रतिनिधिधिकार कार्यालय, भारत सरकार | Copyright Office, Government Of India

पंजीकरण संख्या/Registration Number: **L-147405/2024**

2. आवेदक का नाम, पता तथा राष्ट्रीयता
Name, address and nationality of the applicant

3. कृति के प्रतिनिधिधिकार में आवेदक के हित की प्रकृति
Nature of the applicant's interest in the copyright of the work

4. कृति का वर्ग और वर्ण
Class and description of the work

5. कृति का शीर्षक
Title of the work

LITERARY/ DRAMATIC WORK THIS WORK OUTLINES A CLINICAL RESEARCH AND EXPERIENCE BASED PROTOCOL TO MANAGE THE PATIENT WITH BELL'S PALSY OR IDIOPATHIC LOWER MOTOR FACIAL PALSY.

DARSH GUNJAN'S BELL'S PALSY TREATMENT PROTOCOL

Darsh Gunjan's Bell's palsy treatment protocol

Steps to approach a patient with confirmed Idiopathic lower motor facial palsy (Bell's Palsy) in Neuro-rehabilitation department

Step 1	Baseline evaluation through videography, House Brackmann scale, Sunny Brooke's facial grading scale, Facial nerve conduction study.
Step 2	Protocol based on baseline evaluation will include: - Facial Neuromuscular exercises (for 10 minutes) - Facial massage (for 5 minutes with hypoallergic face oil as per patient's skin type) - Facial therapeutic-ultrasound (at evidence based dosage: frequency: 1000 kHz, intensity: 0.5w/cm2, on-off ratio: 1:2, duration 5 min, over mastoid process, painful area identified behind the ear and paralysed muscles delivered through ELECTROSON 900 by Technomed Electronics) - LOW LEVEL LASER therapy (at evidence based dosage: wavelength of 980 nm and frequency of 100 Hz, energy density of 5 J per every point of direct contact with the skin of the face over the superficial nerve courses of the facial nerve, delivered through Tech Laser 302) - Advanced Pneumatic Radial shockwave therapy Extracorporeal shockwave therapy (at evidence based dosage: frequency 2 Hz, Intensity 2 bars, shots: 500, TD-15 delivered through Shockwave pro 1000 by Technomed Electronics) based on degenerative index as calculated from facial nerve conduction study - Facial taping (to be removed while sleeping)/ Facial splint - Eye care as appropriate
Step 3	Patient will be asked to perform facial neuromuscular exercises 3 times a day followed by facial massage at home
Step 4	Except LLLASER & Advanced Pneumatic Radial shockwave therapy Extracorporeal shockwave therapy, which will be given every third day, rest of the therapeutic protocol will be followed every day.
Step 5	Every 21 days patient will be evaluated through videography, House Brackmann scale, Sunny Brooke's



**COPYRIGHT OFFICE
NEW DELHI**
Reg. No. - L-147405/2024
Date 02/05/2024

(Signature)
अधीक्षक

Step 1:

	facial grading scale, Facial nerve conduction study till complete recovery.
--	---

Clinical visit routine

WEEK1-Day 1: After Step one evaluation patient will be given Facial Neuromuscular exercises, Facial Massage, Facial Ultrasound, LLLaser therapy, Facial splint and eye care advice and home protocol advice

WEEK 1- DAY 2: Facial Neuromuscular exercises (for 10 minutes), Facial Massage, Facial Ultrasound

WEEK 1- DAY 3 : Facial Neuromuscular exercises (for 10 minutes), Facial Massage(for 5 minutes), Advanced Pneumatic Radial shockwave therapy Extracorporeal shockwave therapy on facial nerve trunk in front of ear.

COPYRIGHT OFFICE
NEW DELHI
Reg. No. 147405/2023
Date 02/03/2024

WEEK 1- DAY 4: Facial Neuromuscular exercises (for 10 minutes), Facial Massage (for 5 minutes), Facial Ultrasound.

WEEK 1- DAY 5: Facial Neuromuscular exercises (for 10 minutes), Facial Massage (for 5 minutes), Facial Ultrasound.

WEEK 1- DAY 6: Facial Neuromuscular exercises (for 10 minutes), Facial Massage (for 5 minutes), Facial Ultrasound, LLLaser therapy

WEEK 1- DAY 7: Break Day, only home protocol to be followed

WEEK2 – Same routine to be followed as week 1

WEEK 3 – Same routine to be followed as week 1

WEEK 4- STEP 5 undertaken and if needed same routine will be followed as in week 1.

During the course of treatment, if clinically patient shows complete recovery, then Step 5 is repeated and the patient is discharged.

Baseline Evaluation

Accurate baseline assessment forms the foundation of the protocol. Each patient undergoes a comprehensive evaluation comprising:

- **Standardized videographic documentation** of facial movements at rest and during voluntary expressions
- **House–Brackmann Scale (HBS)** for global facial nerve function

- **Sunnybrook Facial Grading System (SBFGS)** for regional and functional assessment
- **Facial nerve conduction study / Electroneurography (ENoG)** to quantify axonal degeneration

This combined assessment ensures both **qualitative and quantitative profiling** of facial nerve dysfunction.

Step 2: Therapeutic Intervention

Based on baseline severity and ENoG findings, patients are enrolled into a structured multimodal intervention program.

Core Therapeutic Components

- Neuromuscular facial exercises
- Manual facial massage and soft tissue mobilization
- Therapeutic ultrasound
- Low-Level Laser Therapy (LLLT)
- Pneumatic / acoustic shockwave therapy
- Facial taping or splinting where indicated
- Structured eye care protocol

Each modality is selected and dosed according to the degree of nerve degeneration and clinical presentation.

Treatment Frequency

- **Daily therapy sessions** for neuromuscular exercises, massage, ultrasound, and eye care
- **LLLT and shockwave therapy** administered **once every third day**

This frequency balances adequate neural stimulation with prevention of fatigue and aberrant motor learning.

Home-Based Rehabilitation Protocol

A critical component of the protocol is **patient-led home rehabilitation**. Patients are instructed to:

- Perform prescribed facial exercises **three times daily**
- Practice gentle facial massage as taught in the clinic
- Follow strict eye protection measures

Home adherence reinforces clinic-based therapy and accelerates cortical and peripheral reorganization.

Monitoring and Follow-Up

Progress is monitored at **21-day intervals**, using the same standardized tools as baseline evaluation:

- Videographic analysis
- House–Brackmann grading
- Sunnybrook Facial Grading System
- Repeat facial nerve conduction studies where indicated

This allows objective tracking of recovery trends and timely modification of the treatment plan.

Observed Recovery Patterns

Based on longitudinal clinical observation in over 200 patients:

- **Weeks 1–3:**
 - Noticeable reduction in facial asymmetry
 - Improved voluntary facial control
- **Weeks 4–6:**
 - Near-complete recovery in most patients
 - Minimal to absent synkinesis

These timelines are consistently shorter than those reported in conventional rehabilitation approaches.

Clinical Effectiveness and Data Overview

Between **2022 and 2025**, over **200 patients** with Bell’s palsy were managed using the Darsh–Gunjan Protocol©.

Key Observations

- Uniform and accelerated recovery across severity grades
- Rapid restoration of facial symmetry and functional expressions
- **Very low incidence of synkinesis and contractures**
- High patient compliance and satisfaction

The outcomes highlight the value of **severity-based physiotherapy guided by objective diagnostics**.

Determinants of Successful Outcomes

The following factors were identified as critical to success:

- **Early diagnosis using ENoG**
- Severity-specific, standardized physiotherapy
- Avoidance of indiscriminate electrical stimulation
- Strong patient adherence to home protocol
- Regular, objective monitoring

ENoG Versus Clinical Grading: Predictive Value

While clinical grading systems such as HBS and SBFGS provide functional assessment, they remain largely **qualitative**. In contrast, **ENoG offers objective, quantitative data** regarding axonal integrity.

The protocol demonstrates that **ENoG-based stratification allows more precise treatment planning**, particularly in moderate to severe facial palsy.

Electrical Stimulation and Synkinesis: A Cautionary Note

Improper or premature electrical stimulation—especially without assessing nerve degeneration—can promote **aberrant reinnervation**, significantly increasing the risk of synkinesis.

The Darsh–Gunjan Protocol© emphasizes **selective, judicious use of stimulation**, only when nerve integrity and recovery phase permit.

Future Directions

Ongoing and future advancements include:

- Refinement of ENoG interpretation protocols
- Integration of **AI-based facial analysis and diagnostics**
- Large-scale, multicenter clinical studies
- Expansion into **tele-rehabilitation and remote monitoring platforms**

These developments aim to further standardize and globalize facial palsy rehabilitation.

Key Conclusion

The Darsh–Gunjan’s Bell’s Palsy Treatment Protocol© demonstrates that **integrating objective ENoG data with tailored, multimodal physiotherapy** leads to **faster, near-complete recovery** while minimizing long-term complications such as synkinesis and contractures. This protocol represents a reproducible, evidence-informed model for modern facial palsy rehabilitation.

CHAPTER 11

Multidisciplinary Approach and Future Directions in Facial Palsy Management

Dr. Darshpreet Kaur (PT)

Dr. (Prof) Gunjan Kumar

Introduction

Bell's palsy is a multifaceted neurological condition that extends beyond isolated facial weakness. Optimal management requires coordinated efforts across medical, diagnostic, rehabilitative, and psychosocial domains. Increasing evidence supports a **multidisciplinary, patient-centered approach** as the most effective strategy for achieving early recovery, minimizing complications, and improving quality of life.

In parallel, advances in artificial intelligence, digital health platforms, and remote care delivery are reshaping the future of facial palsy management. This chapter outlines the structure of a multidisciplinary care model and explores emerging directions that will define next-generation rehabilitation.

Multidisciplinary Team in Bell's Palsy Management

A comprehensive multidisciplinary team for Bell's palsy typically includes:

- **Neurologist** – Responsible for clinical diagnosis, exclusion of central causes, and pharmacological management
- **Otolaryngologist (ENT Specialist)** – Evaluates temporal bone pathology, middle ear disease, and surgical indications
- **Radiologist** – Provides imaging-based localization and prognostic insight through MRI, HRCT, and ultrasonography
- **Electrophysiologist** – Conducts and interprets ENoG, EMG, and blink reflex studies

- **Physiotherapist** – Designs and executes stage-specific facial rehabilitation and neuromuscular re-education

Collaboration among these specialists ensures accurate diagnosis, objective severity assessment, timely intervention, and structured rehabilitation, resulting in **superior functional outcomes** compared to isolated care.

Role of Artificial Intelligence in Facial Palsy

AI-Assisted Diagnosis and Classification

Artificial intelligence (AI) has significant potential in the early detection and classification of facial nerve dysfunction. Using computer vision and machine learning algorithms, AI systems can analyze facial images or videos to:

- Detect subtle asymmetries at rest and during movement
- Quantify muscle weakness and facial excursion
- Identify early synkinesis
- Objectively grade severity using validated scales

AI-assisted analysis can also help differentiate **central versus peripheral facial palsy**, improving diagnostic accuracy, especially in early or ambiguous cases.

AI in Electrophysiological and Imaging Analysis

AI-driven tools can process large datasets from EMG, ENoG, and imaging studies to:

- Identify patterns predictive of recovery
- Assist in severity stratification
- Support clinical decision-making

These tools hold promise for reducing inter-observer variability and enhancing precision medicine.

Tele-Rehabilitation in Bell's Palsy

Core Components of an Effective Tele-Rehab Protocol

An effective tele-rehabilitation model for Bell's palsy includes:

- Initial **virtual clinical assessment** and grading
- Customized **home exercise programs** with video demonstrations
- Scheduled **live video sessions** with physiotherapists
- Use of **mobile applications or wearable devices** for facial motion tracking
- Structured follow-up with feedback and progression

Tele-rehab protocols must be evidence-based, supervised, and integrated with in-person care when required.

Impact on Accessibility and Equity

Tele-rehabilitation significantly improves access to care, particularly for:

- Patients in rural or underserved regions
- Individuals with mobility limitations
- Those unable to attend frequent in-person sessions
- Situations such as pandemics or travel restrictions

By leveraging smartphones and internet connectivity, tele-rehab promotes **equitable, continuous, and patient-centered care**.

Integrative Care Models in Facial Palsy

Practical Implementation

An integrative care model combines:

- **Conventional medical therapy** (e.g., corticosteroids, antivirals)
- **Structured physiotherapy and rehabilitation**

- **Psychological counseling** to address anxiety, self-image, and social withdrawal
- **Complementary therapies** such as acupuncture, yoga, or relaxation techniques

Care is individualized, goal-oriented, and delivered through shared decision-making between the patient and the multidisciplinary team.

Benefits of Integrative Care

- Holistic management of physical and emotional aspects
- Improved patient adherence and satisfaction
- Enhanced functional and psychosocial outcomes

Integrative models recognize facial palsy as both a **neuromuscular and psychosocial condition**, requiring comprehensive care.

Future Directions in Bell's Palsy Management

The future of facial palsy care lies in **precision rehabilitation**, driven by technology and collaborative models. Key directions include:

- Refinement of **AI-based diagnostic and grading tools**
- Expansion of **tele-rehabilitation platforms** with real-time feedback
- Integration of imaging, electrodiagnostics, and AI into unified dashboards
- Large-scale, multicenter outcome studies
- Development of predictive algorithms for individualized recovery timelines

These advancements aim to standardize care while allowing personalized treatment strategies.

Conclusion: The management of Bell's palsy is evolving from isolated, discipline-specific interventions toward **integrated, technology-enabled care models**. A coordinated multidisciplinary approach—supported by objective diagnostics, advanced rehabilitation techniques, and emerging digital tools—offers the greatest potential for consistent, high-quality outcomes.

As artificial intelligence, tele-rehabilitation, and data-driven protocols continue to mature, the future of facial palsy rehabilitation lies in **precision, accessibility, and collaborative care.**

Further Reading:

Grossman A, Ulanovski D, Barenboim E, Azaria B, Goldstein L. Facial nerve palsy aboard a commercial aircraft. *Aviat Space Environ Med.* 2004 Dec;75(12):1075-6.

Banks CA, Bhama PK, Park J, Hadlock CR, Hadlock TA. Clinician-Graded Electronic Facial Paralysis Assessment: The eFACE. *Plast Reconstr Surg.* 2015 Aug;136(2):223e-230e.

Ojha PT, Nagendra S, Ansari A, Kadam ND, Mathur A, Gopinathan N, Ojha N, Patel H, Bansode A, Kalel O, Morwani KP, Jagtap D, Kumar S, Vij V. Validation of a New Graphic Facial Nerve Grading System: FAME Scale. *Ann Indian Acad Neurol.* 2022 May-Jun;25(3):464-472.

Coulson SE, Croxson GR, Adams RD, O'Dwyer NJ. Reliability of the "Sydney," "Sunnybrook," and "House Brackmann" facial grading systems to assess voluntary movement and synkinesis after facial nerve paralysis. *Otolaryngol Head Neck Surg.* 2005 Apr;132(4):543-9.

Butler DP, Grobbelaar AO. Facial palsy: what can the multidisciplinary team do? *J Multidiscip Healthc.* 2017;10:377-381.

Hadlock TA, Greenfield LJ, Wernick-Robinson M, Cheney ML. Multimodality approach to management of the paralyzed face. *Laryngoscope.* 2006 Aug;116(8):1385-9

Glass GE, Tzafetta K. Bell's palsy: a summary of current evidence and referral algorithm. *Fam Pract.* 2014 Dec;31(6):631-42

CHAPTER 12

Evidence Based Practices

Dr. Darshpreet Kaur (PT)

Dr. Vidhi Singh (PT)

Dr. (Prof) Sneha V. Ketke

1. Acute Peripheral Facial Palsy: Recent Guidelines and Systematic Review (Kim & Lee, 2020)

Bell's palsy is the most common cause of acute peripheral facial palsy. Early **oral corticosteroids** significantly improve recovery; antivirals provide limited additional benefit except in severe cases or Ramsay Hunt syndrome. Eye protection is essential.

2. Bilateral Facial Palsy (Jung et al., 2019)

Bilateral facial palsy is rare and usually indicates **systemic disease** rather than idiopathic Bell's palsy. Common causes include **Guillain–Barré syndrome, infections, autoimmune disorders, and trauma**. Thorough systemic evaluation is mandatory.

3. Facial Exercise Therapy for Facial Palsy (Pereira et al., 2011)

Facial exercise therapy shows **moderate benefit** in improving facial symmetry and motor function, particularly in chronic facial palsy. Evidence supports its use as an adjunct, though optimal protocols remain unclear.

4. Facial Rehabilitation: Evaluation and Treatment Strategies (Robinson & Baiungo, 2018)

Highlights standardized outcome measures and individualized rehabilitation strategies for acute and chronic facial palsy. Emphasizes **neuromuscular retraining**, biofeedback, and post-reanimation therapy.

5. Acquired Bilateral Facial Palsy: Systematic Review (Molinari et al., 2023)

Most bilateral facial palsy cases are secondary, with **GBS, Lyme disease, sarcoidosis, and infections** being common etiologies. Management depends on identifying and treating the underlying cause rather than symptomatic therapy alone.

6. Facial Palsy (Cavoy, 2013)

Stresses the importance of distinguishing **central vs peripheral facial palsy**. Peripheral palsy affects the entire face, whereas central lesions spare the forehead. Early recognition guides appropriate investigations.

7. Facial Nerve Palsy: Clinical Practice and Cognitive Errors (George et al., 2020)

Warns against mislabeling facial palsy as Bell's palsy without adequate evaluation. Encourages the term **“idiopathic facial nerve paralysis”** and highlights common diagnostic pitfalls.

8. Botulinum Toxin Type A in Facial Palsy (de Sanctis Pecora & Shitara, 2021)

Botulinum toxin improves **facial symmetry, synkinesis, and quality of life** in chronic facial palsy. Provides practical guidance on indications, dosing, and injection patterns.

9. Emotional Facial Palsy (Miele et al., 2022)

Describes **automatic–voluntary dissociation**, where voluntary facial movements are impaired but emotional expressions are preserved. Typically associated with **supranuclear (cortical or subcortical) lesions**.

10. Surgical Management of Acute Facial Palsy (Sun et al., 2018)

Most Bell's palsy cases recover medically; surgery is reserved for **severe traumatic palsy or confirmed nerve degeneration**. Timely intervention is crucial for optimal outcomes.

11. Idiopathic Facial Palsy: Umbrella Review (Agostini et al., 2020)

Rehabilitation techniques such as **facial exercises and neuromuscular retraining** show benefit. Evidence for electrotherapy is weak or inconsistent.

12. A General Approach to Facial Palsy (Jowett, 2018)

Proposes a structured framework based on **etiology, timing, severity, and prognosis**. Emphasizes multidisciplinary management and patient-specific treatment planning.

13. Congenital Facial Palsy (Zhu & Han, 2019)

Congenital facial palsy may be unilateral or bilateral and arises from **developmental, genetic, or birth-related causes**. Clinical features and prognosis vary by etiology.

14. Benign Parotid Mass and Facial Palsy (Stewart et al., 2021)

Although facial palsy with parotid mass suggests malignancy, **benign causes do exist** (e.g., infection, inflammation). Careful evaluation is required before definitive conclusions.

15. Facial Palsy: Diagnostic Traps (Schroeter et al., 2023)

Highlights common diagnostic errors, especially overlooking **secondary causes** such as infection, vascular disease, or autoimmune conditions. Persistent or atypical palsy warrants further work-up.

16. Etiology, Diagnosis, and Management of Facial Palsy (2000 patients) (Hohman & Hadlock, 2014)

Bell's palsy remains the most frequent diagnosis. Early referral and standardized assessment improve functional and cosmetic outcomes.

17. Developmental Unilateral Facial Palsy in Newborns (Decraene et al., 2020)

Reviews isolated congenital facial palsy presenting at birth. Multidisciplinary evaluation is important to exclude syndromic associations and guide long-term management.

18. Imaging of Alternobaric Facial Palsy (Lovin et al., 2023)

Suggests pressure changes in a hyperpneumatized temporal bone as a possible mechanism for alternobaric facial palsy. Imaging plays a key role in diagnosis.

19. Facial Palsy (Robin, 1991)

Early overview emphasizing clinical recognition and management principles of facial palsy.

20. Therapeutic Management of Traumatic Facial Palsy (Ottavi et al., 2025)

Trauma accounts for a minority of facial palsy cases. Most recover conservatively, but **surgical intervention** is indicated in selected patients based on severity and nerve integrity.

21. Artificial Intelligence in Facial Palsy Treatment (Müller et al., 2025)

AI-based tools are increasingly used for **objective facial assessment, grading severity, and monitoring recovery** in facial palsy. Machine learning improves accuracy and reproducibility compared to clinician-based scales. The authors recommend standardized datasets and clinical integration.

22. Facial Nerve Palsy: Anatomy, Etiology, Evaluation, and Management (Mavrikakis, 2008)

Provides a comprehensive overview for ophthalmologists, emphasizing **early eye protection**. Stresses identifying underlying causes and

differentiating peripheral from central palsy to prevent corneal complications.

23. Peripheral Facial Nerve Palsy (Pons et al., 2013)

Peripheral facial palsy involves both upper and lower face, unlike central palsy. Bell's palsy is the most common cause; management includes **corticosteroids, eye care, and etiologic evaluation**.

24. Differential Diagnosis of Facial Nerve Palsy (James, 1997)

Highlights **sarcoidosis** as a key cause of bilateral facial palsy in young adults. Facial palsy may be associated with parotid enlargement and uveitis, and often resolves with treatment of the systemic disease.

25. Facial Nerve Rehabilitation (Ishii, 2016)

Reviews principles of facial nerve rehabilitation across age groups. Emphasizes **early therapy, prevention of synkinesis**, and long-term functional optimization.

26. Pediatric Facial Reanimation (Hadford et al., 2024)

Pediatric facial palsy causes significant developmental and psychosocial impact. Treatment focuses on **cause-specific management** and dynamic reanimation procedures tailored to growth and neuroplasticity.

27. Postradiotherapy Facial Palsy (Sbai et al., 2020)

Facial palsy following radiotherapy raises concern for **radiation-induced neuropathy, tumor recurrence, or infection**. Careful diagnostic evaluation is required.

28. Facial Palsy: A Leprosy Surgeon's Viewpoint (Warren, 1990)

Facial palsy in leprosy threatens vision and facial symmetry. Surgical techniques using **temporal muscle and fascia** can restore eyelid closure and oral competence.

29. Facial Palsy after Orthognathic Surgery (Hattori et al., 2024)

Facial palsy is a rare complication ($\approx 0.13\%$) after orthognathic surgery. Most cases recover spontaneously; injury is usually due to **traction or compression** rather than nerve transection.

30. Facial Palsy (Ludman, 1981)

Classic overview of causes, presentation, and management of facial palsy, emphasizing early diagnosis and conservative treatment.

31. Differential Diagnosis of Facial Nerve Palsy (May & Klein, 1991)

Comprehensive review of conditions causing facial palsy. Strongly emphasizes identifying secondary causes to guide appropriate treatment.

32. Recurrent Facial Palsy and Fissured Tongue (Sharma & Sharma, 2021)

Suggests **Melkersson–Rosenthal syndrome** in patients with recurrent facial palsy and fissured tongue. Episodes often resolve completely.

33. Cranial Nerve Defects in Congenital Facial Palsy (Carr et al., 1997)

Congenital facial palsy may be isolated or associated with other cranial nerve deficits. Bilateral involvement is more common in females.

34. Emotional Impact of Facial Palsy (Dobel et al., 2013)

Facial palsy significantly affects **emotional expression, communication, and quality of life**, even in the acute phase. Psychological support is an important part of management.

35. Childhood Peripheral Facial Palsy (Karalok et al., 2018)

Most pediatric cases are idiopathic with good prognosis. Corticosteroid treatment is commonly used, and recurrence is uncommon but possible.

36. Delayed Facial Palsy after Cochlear Implantation (Ishikawa et al., 2022)

Delayed facial palsy post-cochlear implantation may result from **herpesvirus reactivation**. Antiviral therapy leads to good recovery in most cases.

37. Post-COVID-19 Bilateral Facial Nerve Palsy (Inui et al., 2022)

Reports rare bilateral facial palsy occurring weeks after COVID-19 infection. Suggests an **immune-mediated mechanism**.

38. Idiopathic Recurrent Facial Palsy (Scola Yurrita et al., 2004)

Idiopathic recurrent facial palsy is rare and typically shows **complete recovery between episodes**. Diagnosis is one of exclusion.

39. Facial Palsy after Embolization with Squid 12 (Moreno-Paredes et al., 2021)

Facial palsy can occur as a rare complication of endovascular embolization. Imaging may be normal, and recovery is often favorable.

40. Facial Palsy (Cawthorne & Haynes, 1956)

Early foundational article outlining clinical features and treatment principles of facial palsy.

41. Facial Palsy in COVID-19 Patient (Paudel et al., 2022)

COVID-19–associated facial palsy is rare but reported. Most cases resemble Bell’s palsy and recover with standard treatment.

42. Management of Peripheral Facial Nerve Palsy in Children (Tabarki, 2014)

Idiopathic facial palsy is the most common pediatric form. Diagnosis is by exclusion, and prognosis is generally excellent.

43. Facial Palsy and PRES due to Lyme Disease (Okumura et al., 2022)

Illustrates facial palsy as an early manifestation of **Lyme disease**, with rare progression to PRES. Highlights importance of infectious evaluation.

44. Facial Palsy in Children: Long-Term Outcome (Kanerva et al., 2021)

Most children recover well, but **recurrence rates are higher than expected**. Long-term follow-up is recommended.

45. Childhood Facial Palsy: Etiologic Factors (Hanci et al., 2019)

Idiopathic palsy is most common; viral infections are frequent secondary causes. Early corticosteroid therapy is associated with better outcomes.

46. Prevention and Rehabilitation of Facial Palsy in Vestibular Schwannoma (Freeman et al., 2025)

Facial nerve outcomes depend on **extent of nerve injury and recovery time**. Rehabilitation and reanimation procedures are used when recovery is incomplete.

47. Etiology and Management of Facial Palsy (Boerner & Seiff, 1994)

Reviews facial nerve anatomy and consequences of proximal lesions (e.g., loss of lacrimation, taste). Emphasizes individualized management.

48. Facial Nerve Pathology in Children (Derise et al., 2024)

Congenital causes are more common in children than adults. Management prioritizes **dynamic restoration of facial expression**.

49. Assessment of Eye Closure and Blink in Facial Palsy (Zaidman et al., 2021)

Reviews tools for assessing eye closure and blink, critical for ocular protection. Highlights lack of standardized, validated assessment systems

51. Quality of Life after Surgical Treatments for Facial Palsy (van der Lely et al., 2024)

Surgical facial reanimation significantly improves **quality of life (QoL)**, especially emotional and social domains. Dynamic procedures show greater QoL gains than static ones. PROMs are essential for outcome assessment.

52. A Surgical Algorithm for the Management of Facial Palsy (Mackinnon & Dellon, 1988)

Classic algorithm based on **timing, nerve continuity, and muscle viability**. Early nerve repair is preferred; delayed cases require muscle or tendon transfer.

53. Peripheral Facial Palsy in Children (Yilmaz et al., 2014)

Most pediatric facial palsy cases are **idiopathic** with good prognosis. Infections and trauma are common secondary causes. Recovery is generally better than in adults.

54. Facial Palsy-Specific Quality of Life in 920 Patients (Tavares-Brito et al., 2019)

QoL correlates only **moderately** with clinician-graded severity. Psychosocial factors and synkinesis significantly influence patient-reported outcomes.

55. Ophthalmic Assessment and Management in Acute Facial Palsy (Malik et al., 2025)

Highlights the critical role of **early eye protection** to prevent exposure keratopathy. Management should be individualized based on blink, lagophthalmos, and tear function.

56. Viral Infection and Immune Response in Facial Palsy Pathogenesis (Wang et al., 2025)

Facial palsy results from a **synergistic interaction between viral infection and host immune response**. HSV-1, VZV, and SARS-CoV-2 play key roles via inflammatory nerve damage.

57. Facial Reconstruction and Rehabilitation (Guntinas-Lichius et al., 2016)

Malignancy-related facial palsy requires **oncologic safety first**, followed by reconstruction. Combines nerve repair, reanimation, and rehabilitation.

58. Acute Peripheral Facial Palsy in Older Adults (Jeong et al., 2022)

Older patients show higher rates of **secondary causes** and slower recovery. Comorbidities influence prognosis more than in younger patients.

59. Facial Palsy after Dental Procedures (Gaudin et al., 2017)

Delayed facial palsy after dental work is rare ($\approx 0.7\%$). Most cases resemble **Bell's palsy**, likely due to viral reactivation rather than direct nerve injury.

60. Functional Results after Facial Reanimation in Iatrogenic Facial Palsy (Györi et al., 2020)

Facial reanimation improves symmetry and function in iatrogenic palsy. Outcomes depend on **patient age, timing, and technique** used.

61. Beyond Botox: Management of Nonflaccid Facial Palsy (Miller & Hadlock, 2020)

Nonflaccid facial palsy is characterized by **synkinesis and hypertonicity**. Management includes botulinum toxin, neuromuscular retraining, and selective neurectomy.

62. Reanimation following Facial Palsy: Present and Future Directions (Ramakrishnan et al., 2010)

Advocates a **multidisciplinary, tailored approach** combining surgery, physiotherapy, and adjunctive treatments for chronic facial palsy.

63. Facial Nerve Palsy in Lyme Neuroborreliosis (Marques et al., 2022)

Facial palsy is the most common manifestation of Lyme neuroborreliosis, often **bilateral or sequential**. Prognosis is excellent with antibiotic therapy.

64. Free Gracilis Transfer and Static Suspension (Jowett & Hadlock, 2018)

Free gracilis muscle transfer is the **gold standard** for long-standing flaccid facial palsy. Static suspension improves symmetry at rest.

65. Facial Palsy – When Facial Expression Disappears (Berghaus & San Nicoló, 2015)

Reviews functional, emotional, and social consequences of facial palsy. Emphasizes holistic management beyond motor recovery.

66. Facial Palsy, Bilateral (Williams, 1949)

Early recognition that bilateral facial palsy usually reflects **systemic disease**, not idiopathic Bell's palsy.

67. Brazilian Society of Otology Guidelines on Facial Palsy (Pauna et al., 2024)

Comprehensive guideline covering **diagnosis, electrophysiology, and management** of idiopathic, traumatic, recurrent, and Ramsay Hunt-related facial palsy.

68. Facial Paralysis (Gates, 1987)

Uses the term **idiopathic facial paralysis** instead of Bell's palsy. Reviews nontraumatic causes and conservative management.

69. Facial Palsy in Dental Surgery (Miles, 1992)

Dental procedures can rarely precipitate facial palsy via **local anesthesia, trauma, or viral reactivation**.

70. Facial Palsy in Kawasaki Disease (Hattori et al., 1987)

Facial palsy is a rare neurological manifestation of Kawasaki disease, likely related to **vasculitis**. Usually transient.

71. Facial Palsy (Bateman, 1992)

Provides a practical diagnostic approach based on **lesion localization** and associated neurological features.

72. Facial Palsy Induced by Covaxin (Chaudhari et al., 2023)

Reports a rare case of post-vaccination facial palsy. Course and recovery resemble idiopathic Bell's palsy.

73. Facial Palsy in Tetanus (Achari, 1973)

Facial palsy may occur in tetanus due to neurotoxin-mediated cranial nerve involvement.

74. Traumatic Facial Nerve Palsy: Surgical vs Nonsurgical Outcomes (Bae et al., 2023)

Immediate, complete traumatic palsy may benefit from decompression surgery. Partial palsy often recovers well with conservative management.

75. Current Concepts for Eyelid Reanimation (Momeni & Khosla, 2014)

Eyelid reanimation is essential to protect vision. Techniques include **gold weight implantation, tarsorrhaphy, and dynamic reanimation.**

76. MEEI Facial Palsy Photo and Video Standard Set (Greene et al., 2020)

Introduces standardized photo/video datasets and AI tools to improve **objective facial palsy assessment.**

77. Delayed Facial Palsy after Anterior Petrosal Approach (Guthikonda et al., 2010)

Delayed palsy can occur without direct nerve injury, likely due to **viral reactivation or inflammation.** Responds to antivirals and steroids.

78. Prognostic Accuracy in Acute Peripheral Facial Palsy (Aoyagi, 2005)

Prognosis depends mainly on **cause, site, and severity of nerve injury.** Early electrophysiologic testing improves prediction.

79. Approach to Facial Weakness (Wang et al., 2021)

Provides a structured neurologic approach distinguishing **central vs peripheral facial weakness,** with emphasis on anatomy and systemic causes.

About the Authors

Dr. Darshpreet Kaur (PT) and **Dr. (Prof) Gunjan Kumar** are a dedicated clinician–researcher team whose collaborative work has significantly advanced the field of facial nerve rehabilitation in India.

Dr. Darshpreet Kaur (PT) is an expert in rehabilitation sciences with specialized experience in neuro-rehabilitation and evidence-based physiotherapy. Dr. Gunjan Kumar is a practicing neurologist with extensive clinical expertise in the diagnosis and management of neurological disorders. Together, they lead one of Bihar’s most advanced evidence-based physiotherapy and neurology centers—**Neuro Vihar**, a hub recognized for its integrated, patient-centric approach to rehabilitation. Neuro Vihar has expanded its expert care from Patna to Danapur @ Jeevandan Health and Jalandhar, Punjab.

Their close clinical collaboration intensified during the COVID-19 pandemic, when they encountered a growing number of Bell’s palsy patients presenting with **synkinesis, delayed recovery, and long-term functional deficits** following conventional and often outdated therapeutic practices. These observations prompted them to critically evaluate existing treatment paradigms and challenge long-standing assumptions in facial palsy management.

Driven by a shared commitment to clinical excellence, they developed a **novel, multidimensional rehabilitation approach** grounded in neurological precision, objective diagnostics, and functional physiotherapy. This work culminated in the creation of **Darsh–Gunjan’s Bell’s Palsy Treatment Protocol©**—a structured, scientifically grounded methodology that became **India’s first government-copyrighted protocol dedicated exclusively to facial nerve rehabilitation**.

Through this protocol, they have transformed recovery outcomes for hundreds of patients, achieving faster, more complete functional restoration with a markedly reduced incidence of long-term complications such as synkinesis and facial contractures. Their work

continues to influence clinical practice and is paving the way for a broader national shift toward **precision-based, protocol-driven management of Bell's palsy**.

Their shared vision remains focused on restoring not only facial movement, but also symmetry, confidence, and quality of life for every patient they serve.

“Helping faces recover is our shared purpose.”